

This is a repository copy of *Shining a light on the Arabidopsis circadian clock*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/119697/>

Version: Accepted Version

Article:

Oakenfull, Rachael and Davis, Seth Jon orcid.org/0000-0001-5928-9046 (2017) Shining a light on the Arabidopsis circadian clock. *Plant, Cell and Environment*. 2571–2585. ISSN 0140-7791

<https://doi.org/10.1111/pce.13033>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Shining a light on the Arabidopsis circadian clock

Journal:	<i>Plant, Cell & Environment</i>
Manuscript ID	PCE-17-0403.R1
Wiley - Manuscript type:	Invited Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Oakenfull, Rachael; University of York, Department of Biology Davis, Seth Jon; University of York, Department of Biology
Environment Keywords:	circadian, light quality
Physiology Keywords:	signaling
Other Keywords:	
Abstract:	<p>The circadian clock provides essential timing information to ensure optimal growth to prevailing external environmental conditions. A major time-setting mechanism (zeitgeber) in clock synchronisation is light. Differing light wavelengths, intensities and photoperiodic duration are processed for the clock-setting mechanism. Many studies on light-input pathways to the clock have focused on <i>Arabidopsis thaliana</i>. Photoreceptors are specific chromic proteins that detect light signals and transmit this information to the central circadian oscillator through a number of different signalling mechanisms. The most well characterised clock-mediating photoreceptors are cryptochromes and phytochromes, detecting blue, red and far-red wavelengths of light. Ultraviolet and shaded light are also processed signals to the oscillator. Notably, the clock reciprocally generates rhythms of photoreceptor action leading to so-called gating of light responses. Intermediate proteins, such as Phytochrome interacting factors (PIFs), constitutive photomorphogenic 1 (COP1) and EARLY FLOWERING 3 (ELF3), have been established in signalling pathways downstream of photoreceptor activation. However, the precise details for these signalling mechanisms are not fully established. This review highlights both historical and recent efforts made to understand overall light input to the oscillator, first looking at how each wavelength of light is detected, this is then related to known input mechanisms and their interactions.</p>

For Review Only

Shining a light on the Arabidopsis circadian clock

Rachael J. Oakenfull, Seth J. Davis

Department of Biology, University of York, York, UK

Corresponding author: Seth J. Davis

E-mail: seth.davis@york.ac.uk

For Review Only

14 Abstract

15

16 The circadian clock provides essential timing information to ensure optimal growth to prevailing
17 external environmental conditions. A major time-setting mechanism (*zeitgeber*) in clock
18 synchronisation is light. Differing light wavelengths, intensities and photoperiodic duration are
19 processed for the clock-setting mechanism. Many studies on light-input pathways to the clock have
20 focused on *Arabidopsis thaliana*. Photoreceptors are specific chromic proteins that detect light
21 signals and transmit this information to the central circadian oscillator through a number of different
22 signalling mechanisms. The most well characterised clock-mediating photoreceptors are
23 cryptochromes and phytochromes, detecting blue, red and far-red wavelengths of light. Ultraviolet
24 and shaded light are also processed signals to the oscillator. Notably, the clock reciprocally
25 generates rhythms of photoreceptor action leading to so-called gating of light responses.
26 Intermediate proteins, such as Phytochrome interacting factors (PIFs), constitutive
27 photomorphogenic 1 (COP1) and EARLY FLOWERING 3 (ELF3), have been established in signalling
28 pathways downstream of photoreceptor activation. However, the precise details for these signalling
29 mechanisms are not fully established. This review highlights both historical and recent efforts made
30 to understand overall light input to the oscillator, first looking at how each wavelength of light is
31 detected, this is then related to known input mechanisms and their interactions.

32

33 The circadian clock

34 The circadian clock allows plants as sessile organisms to synchronise with diurnal changes in
 35 the environment (Dodd *et al.*, 2005). Daily external environmental stimuli are required to initiate
 36 circadian oscillations and to maintain synchronicity with the external environment. This process is
 37 called entrainment. The environmental cues governing these processes are termed *zeitgebers* (from
 38 German: "time givers"). The ability to synchronise with the external environment efficiently confers
 39 enhanced fitness (Michael *et al.*, 2003).

40 Diurnal changes in cellular processes controlled by the clock allow plants to anticipate, and
 41 therefore better survive, a range of stresses (Sanchez *et al.*, 2011). Diurnal changes have been
 42 shown to occur in cold/freezing tolerance (Fornara *et al.*, 2015, Nakamichi *et al.*, 2009), drought
 43 tolerance (Habte *et al.*, 2014), pathogen response (Shin *et al.*, 2012, Wang *et al.*, 2011) and
 44 photosynthesis (Pyl *et al.*, 2012). This synchronisation is the product of a large number of
 45 rhythmically regulated cellular processes (Bujdoso & Davis, 2013, Hanano *et al.*, 2008), many of
 46 which are triggered by light perception (Wenden *et al.*, 2011). Perception of daily *zeitgebers*, such as
 47 changes in light and temperature (Chow *et al.*, 2014, Harmer, 2009), enable plants to reset the clock
 48 at dawn, and feed back to the central oscillator. For this light entrainment, photoreceptors play a
 49 major role (Somers *et al.*, 1998a, Toth *et al.*, 2001). The circadian clock was derived from the
 50 principle of inter-connected, positive and negative feedback loops (Shearman *et al.*, 2000). For the
 51 purpose of this review, light input into the *Arabidopsis thaliana* (*Arabidopsis*) circadian clock will be
 52 the focus of discussion, with a brief examination of clock components.

53 In *Arabidopsis*, morning expressed Myb-like transcription factors *CIRCADIAN CLOCK*
 54 *ASSOCIATED 1* (*CCA1*) (Wang & Tobin, 1998) and *LATE ELONGATED HYPOCOTYL* (*LHY*) (Schaffer *et al.*,
 55 1998) antagonize expression of the evening expressed pseudo-response regulator (PRR) *TIMING OF*
 56 *CAB EXPRESSION 1* (*TOC1*) (Strayer *et al.*, 2000). These three genes form the core negative feedback
 57 loop of the circadian oscillator (Alabadí *et al.*, 2001, Gendron *et al.*, 2012) (Figure 1). Several other
 58 genes form additional loops within this core oscillator. In day time *CCA1* and *LHY* repress expression
 59 of the pseudo-response regulators *PRR5*, *PRR7* and *PRR9* (Adams *et al.*, 2015, Kamioka *et al.*, 2016),
 60 as well as *TOC1*, *GI*, and the genes that generate the evening complex (Locke *et al.*, 2006, Nakamichi
 61 *et al.*, 2009, Pokhilko *et al.*, 2010, Zeilinger *et al.*, 2006). *GIGANTEA* (*GI*) is evening expressed and is
 62 proposed to form an additional negative feedback-loop with *TOC1* (Locke *et al.*, 2006). All of these
 63 loops are connected through the action of the evening complex formed by *LUX ARRHYTHMO* (*LUX*),
 64 *EARLY FLOWERING 3* (*ELF3*) and *EARLY FLOWERING 4* (*ELF4*), which directly inhibits the expression of
 65 *PRR9* (Helfer *et al.*, 2011, Herrero *et al.*, 2012), *PRR7*, *GI* and *LUX* (Mizuno *et al.*, 2014). The absence
 66 of even one component of the evening complex gives rise to plants that are photoperiod insensitive.
 67 This results in early flowering, long hypocotyl growth and arrhythmicity of the free-running circadian
 68 period (Hazen *et al.*, 2005, McWatters *et al.*, 2007, Onai & Ishiura, 2005, Thines & Harmon, 2010).
 69 The importance of the three evening-complex components is thusly highlighted in maintaining a
 70 functional circadian clock, and therefore the physiological processes controlled by the clock, such as
 71 the input of diurnal photoperiod information, (Covington *et al.*, 2001, Más *et al.*, 2003, Mizoguchi *et*
 72 *al.*, 2005, Park *et al.*, 1999).

73

74 Effects of light on the clock

75 Light changes throughout a day-night cycle are pronounced and thus robustly entrain the
 76 clock. In the light phase of a daily cycle, the dark to light transition of dawn is used as a time setting
 77 checkpoint (Millar *et al.*, 1995). Prolonged darkness causes many of the core genes in the
 78 *Arabidopsis* central oscillator to rapidly become arrhythmic, due to the lack of essential light time
 79 setting cues (Figure 2a) (Millar *et al.*, 1995). This dampening effect, leading to arrhythmicity is
 80 particularly noticeable in the absence of media containing sucrose. In prolonged darkness, sucrose
 81 can act as a substitute for light in maintaining rhythmicity for a number of days (Bläsing *et al.*, 2005).
 82 Light has two main modes to set the clock. The first is parametric entrainment; gradual entrainment
 83 of the clock, such as the acceleration of the clock induced by increased light perception, which

eventually leads to a phase shift of the clock back to a correct resonance. Parametric entrainment follows Aschoff's rule, as light intensity increases, the speed of the clock increases. As intensity decreases, the speed of the clock slows (Aschoff, 1979), (Figure 2B/C). Increases in light intensity, lead to decreases in periodicity (Somers *et al.*, 1998a). The second light-induced time-setting mechanism is non-parametric entrainment: rapid re-entrainment. This leads to a rapid time setting of the clock at dawn (Millar & Kay, 1996). Non-parametric entrainment requires an extended light exposure far beyond that which activates light-regulated gene expression (Millar & Kay, 1996). Metabolic entrainment is also a mechanism for non-parametric entrainment (Haydon *et al.*, 2013, Sanchez-Villarreal *et al.*, 2013; Haydon & Webb, 2016; Shin *et al.*, 2017). The different photoreceptors and photochromic proteins involved in light entrainment are described in more detail below.

How are different wavelengths of light input to the *Arabidopsis* clock?

Diurnal organisms, particularly plants, are subjected to Aschoff's rule: an increase in light intensity accelerates the circadian-oscillator speed leading to shortening of periodicity (Aschoff, 1979). Light input to the circadian clock is presumed to occur through the action of different types of photoreceptors (Somers *et al.*, 1998a). There are more than ten known circadian-associated photoreceptors (Edwards *et al.*, 2015). These can be split into four classes: phytochromes, cryptochromes, ZTL/FKF1/LKP2 family, and UVR8. Each receptor contributes in the dose-dependent perception of far-red, red, blue, and ultra-violet light (Cashmore *et al.*, 1999, Mas *et al.*, 2000, Rizzini *et al.*, 2011, Song *et al.*, 2012). It is presumed that the input of this information is co-ordinately relayed to the central oscillator.

Both phytochromes and cryptochromes play key roles in light responsive time setting mechanisms, in a manner that follows Aschoff's rule (Devlin & Kay, 2000a, Somers *et al.*, 1998a). This is due to the ability of both phytochromes and cryptochromes to form photoreceptor complexes (Más *et al.*, 2003) that are genetically interactive in clock function (Devlin & Kay, 2000a). Excitation of these photoreceptors cause the central oscillator to accelerate, changing the overall speed of the clock (Devlin & Kay, 2000b, Herrero *et al.*, 2012, Kolmos *et al.*, 2011, Somers *et al.*, 1998a, Somers *et al.*, 1998b). There are a number of different known mechanisms through which light absorption by photoreceptors input environmental information to the oscillator, however, these mechanistic details are not complete. Regulation of transcription by circadian gating restricts changes in RNA levels to specific times of day. Therefore preventing transcription of some light-regulated clock genes in response to unexpected external stimuli, for example light pulses during the night (Millar & Kay, 1996). Light regulation of myb transcription factors, such as *CCA1* and *LHY* effect the transcription and stability of other clock components, such as *PRR9/7* (Carre & Kay, 1995). Messengers such as Ca^{2+} and calmodulin signalling may also affect circadian regulation in response to light (Johnson *et al.*, 1995, Millar & Kay, 1996). Light also directly controls the degradation of *PRR5*, *PRR7*, *PRR9*, *TOC1* and *GI* proteins (Farré and Kay, 2007, Ito *et al.*, 2007, Kiba *et al.*, 2007, Más *et al.*, 2003, Matsushika *et al.*, 2000). These degradation events then act on outputs within a diurnal context, which change in duration throughout the season (Davis, 2002, Guerriero *et al.*, 2012, Salazar *et al.*, 2009, Song *et al.*, 2012, Troein *et al.*, 2009). Light thus has multiple mechanistic inputs to clock processes, all of which control entrainment. How each individual wavelength of light is input to the clock will be discussed below.

Red light

Phytochromes are predominantly red-light photoreceptors, absorbing maximally at wavelengths between 600 and 700nm (Somers *et al.*, 1998a). *Arabidopsis* has five phytochromes (Sharrock & Quail, 1989), phyA-phyE (Mathews & Sharrock, 1997). Each phytochrome acts as a light input sensor to form regulatory feedback loops within the circadian clock. Phytochromes are in turn reported to be negatively regulated by the clock through cryptochrome (CRY) signals (Devlin & Kay, 2000a, Mas *et al.*, 2000). Phytochromes exist in two interconvertible forms; the inactive Pr form is

converted by red light to the active Pfr form which can be converted back to the inactive Pr state by far-red light (Rudiger *et al.*, 1983). These conversion events between active and inactive forms of phytochrome are essential to light input to the clock, as discussed below (see far red, PIFs, ELF3). Each of the five phytochromes play distinct roles in light sensing.

phyA mediates entrainment responses to low intensity red light and pulses of light (Quail *et al.*, 1995, Somers *et al.*, 1998a). A *PHYA* deficiency mutation, results in an altered period length in dim red light (Somers *et al.*, 1998a). It is not known how phyA signals to the clock as it has not been reported to directly bind to a clock-associated factor, in contrast to the other four phytochromes (Huang *et al.*, 2016).

phyB is the main detector for high intensity red light (Somers *et al.*, 1998a). Both phyB and phyD are able to detect red and far-red wavelengths of light (Aukerman *et al.*, 1997, Devlin *et al.*, 1999). phyB physically interacts with ELF3 in the central oscillator to provide a direct light input to the clock (Kolmos *et al.*, 2011, Liu *et al.*, 2001). *phyb* mutants show an altered response to shade avoidance (Smith, 1995), which is also a phenotype of the *elf3* mutant (Huang *et al.*, 2016). phyC to phyE also interact with ELF3 protein (Huang *et al.*, 2016), but this has not yet been connected to the clock (Liu *et al.*, 2001). Under high fluence red light, *phyb* mutants and the phyB overexpressor have a period defects and altered phase (Anderson *et al.*, 1997, Kolmos *et al.*, 2011, Salomé *et al.*, 2002, Somers *et al.*, 1998a). Also altered cryptochrome signalling (see blue light below), phyB and CRY2 physically interact by translocating to the nucleus in red light (Mas *et al.*, 2000), where phyB is then suppressed by CRY2 (Mas *et al.*, 2000). This alters clock performance under white light conditions (red and blue light together) (Devlin & Kay, 2000a).

In non-peer-reviewed work, phyC was found to play a role in white-light input and red-light detection. Mutations in *PHYC* result in a long-period phenotype, which was shown to be temperature dependant, suggesting that phyC inputs not only light information to the clock, but also temperature (Edwards *et al.*, 2015, Franklin *et al.*, 2003, Qin *et al.*, 1997). phyE along with phyD plays a role in controlling the period length of *CAB* gene expression, however, many of the clock effects of phyE and phyD are masked by phyB (Franklin & Quail, 2010). phyE works with phyB and phyD in the regulation of shade avoidance (Devlin *et al.*, 1998). Interestingly, the promoters of *PHYA* and *PHYB* are down-regulated by light, whereas the *PHYC* promoter is upregulated (Tóth *et al.*, 2001), *PHYD* and *PHYE* do not show changes in expression in response to light changes. *PHYB,D,E* mediate high fluence red light input to the clock with *PHYA,B,D,E* acting additively to input red-light information to the clock, as a result the clock runs faster as the detected intensity of red light increases (Devlin & Kay, 2000a). The absence of all five phytochromes results in severally attenuated rhythms, but not a total loss of clock function (Hu *et al.*, 2013). Together, all five phytochromes play differing roles in mediating light-dependant changes in periodicity.

Blue light

Cryptochromes are blue light (492 to 455nm) and UVA photoreceptors present in both plants and animals (Cashmore *et al.*, 1999). The *HY4* locus was found to encode cryptochrome 1 (CRY1). It was identified due to *cry1* (*hy4*) mutants growing with a long-hypocotyl phenotype and being unable to respond to blue light (Ahmad & Cashmore, 1993, Koornneef *et al.*, 1980), *cry1/hy4* plants are also late flowering (Goto *et al.*, 1991, Millar *et al.*, 1995). *cry1* mutants have a long period under blue light (Somers *et al.*, 1998a), suggesting CRY1 acts as a photoreceptor for blue-light entrainment of the clock (Devlin & Kay, 2000a). Overexpression of *CRY1* caused increased sensitivity to blue light and period shortening (Lin *et al.*, 1996, Somers *et al.*, 1998a). CRY1 is a soluble protein when grown in both light and dark conditions in *Arabidopsis* (Lin *et al.*, 1996), CRY1 is more stable than CRY2 and works at higher light intensities (Lin *et al.*, 1998). Cryptochrome 2 (CRY2) can detect low intensity light and is rapidly degraded under blue light (Lin *et al.*, 1998). In light, *CRY2* promoter activity is down-regulated whereas, *CRY1* is upregulated (Tóth *et al.*, 2001). The *cry2* mutation alters sensitivity to photoperiod and flowering in *Arabidopsis*, but does not have a detectable individual effect on circadian rhythm (Devlin & Kay, 2000a, Guo *et al.*, 1998). Overexpression of either *CRY1* or

CRY2 gives rise to a higher blue-light sensitivity under low light conditions than in the individual overexpression lines (Ahmad *et al.*, 1998a). Double mutant *cry1, cry2* plants are rhythmic, suggesting that although CRY1 inputs blue light into the clock CRY1 and 2 are not part of the central oscillator (Devlin & Kay, 2000a). However, CRY1 and CRY2 gene expression oscillates with a circadian rhythm under constant light (Harmer *et al.*, 2000). CRY1 and 2 work together to input information to the clock in a similar way to phyA and B, but at differing light intensities.

Phytochromes are able to absorb low fluence blue light alongside CRY1 for period length control. *phyA* mutants show a period lengthening effect when free run under blue light (Somers *et al.*, 1998a). Without phyA detection of blue light, the input relies on CRY1 alone causing the period to lengthen as the plant detects less light than the actual ambient intensity of irradiation. Conversely, *PHYA* overexpression has been proposed to cause period shortening under blue light, as more blue light is processed as an input than the actual ambient light intensity. Phytochromes thus also work in blue-light signalling to the clock.

High light synergism - (White light)

White light comprises of multiple light wavelengths. As such, interactions between phytochromes and cryptochromes are needed to input this information into the circadian clock. These interactions were found with loss of function mutants for both phytochromes and cryptochromes. In wild-type plants *CAB2* period decreases as light intensity increases, *cry2* mutants were found to be deficient in a white light response as they have a *CAB2* period increase in response to high light (Mas *et al.*, 2000). This period increase was not detected in either red or blue light alone, suggesting that to be active CRY2 needs multiple wavelengths of light simultaneously and phyB (Mas *et al.*, 2000). CRY1 was also found to be required for phyA signalling as *cry1* and *cry2* mutants are unable to detect red light above the fluence range of both phyA and phyB (Devlin & Kay, 2000a). Light induces nuclear compartmentalisation of phytochromes where phyA and phyB directly interact with CRY1 and CRY2 (Mas *et al.*, 2000), the kinase activity of phyA phosphorylates CRY1 and CRY2 (Ahmad *et al.*, 1998b).

Phytochromes and cryptochromes facilitate signal integration of multiple light cues. CRY2 is activated when illuminated by multiple wavelengths of light suggesting it is needed for phytochrome activation (Mas *et al.*, 2000). This is also highlighted as both cryptochromes reach peak RNA expression with a similar expression pattern to the corresponding phytochromes (Toth *et al.*, 2001). It was found that the active Pfr form of phytochrome is needed for CRY2 expression, CRY2 then suppresses *PHYB* expression. However, PfrB is able to override CRY2 signalling to flowering time control via pathways, such as COP1 (see below). PfrB binding to the intermediate SPA1 allows degradation of the COP1-SPA1 complex, which is needed as an intermediate of CRY1/2 induced inhibition of photomorphogenic factors such as HFR and CO (Mas *et al.*, 2000, Sheerin *et al.*, 2015). Additionally, both CRY1 and CRY2 were found to be phosphorylated by the kinase activity of phyA (Ahmad *et al.*, 1998b). It could be considered that phytochromes and cryptochromes work together in the "white light" response, which is a more than the additive effect of plants grown under blue and red light. However, quadruple mutants for *phyA, phyB, cry1* and *cry2* still showed rhythmic leaf movement in response to light-dark cycles suggesting that other photoreceptors must play a role in overall light input to the circadian clock (Yanovsky *et al.*, 2000). However, the exact relationship between phytochromes and cryptochromes is yet to be resolved.

ZTL family; blue-light absorbing with action under red-light and darkness

Zeitlupe (ZTL), also reported as ADAGIO1 (ADO1), links light input by both cryptochromes and phytochromes to the clock (Jarillo *et al.*, 2001, Kim *et al.*, 2007). *ZTL* mutant lines showed altered cotyledon movement and gene expression under different light conditions (Jarillo *et al.*, 2001). Under blue and white light, *ztl* plants have a long period whereas under red light the *ztl* lines were reported to be arrhythmic for *CCR2* expression, cotyledon movement and stem elongation (Jarillo *et al.*, 2001). *ZTL* mutants were found to have a long-period phenotype for *CAB/TOC1* under

red light (Kevei *et al.*, 2006, Kim *et al.*, 2005). ZTL thus is required for the perception of multiple wavelengths of light into the oscillator.

ZTL encodes a protein reported to be a blue-light photoreceptor, as it contains a PAS domain, F box domain linking proteins to a SCF ubiquitination complex, kelch repeats and a light, oxygen or voltage (LOV) domain allowing protein-protein interactions (Mas *et al.*, 2003). Interactions between TOC1 and ZTL were found to occur through these kelch-repeat zones (Kevei *et al.*, 2006). The PAS/ LOV domain were identified as essential for coupling ZTL to red light (Kevei *et al.*, 2006), which was then found to occur through ZTL binding to the C-terminus of PhyB and CRY1 (Kim *et al.*, 2007).

ZTL is constitutively expressed at the RNA level, however, oscillations in ZTL protein levels are seen (Kim *et al.*, 2007). These are proposed to result from the binding of Gigantea (GI) to maintain the stability of ZTL. ZTL protein folding is chaperoned by HSP90 (Kim *et al.*, 2011), GI binds to the ZTL-HSP90 complex to ensure specificity of protein folding (Cha *et al.*, 2017, Kim *et al.*, 2011). Interactions between ZTL and GI are enhanced by blue light through the LOV domain in ZTL (Kim *et al.*, 2007). ZTL controls proteasomal degradation of TOC1 (Más *et al.*, 2003). This ZTL-GI interaction is believed to control a central part of the circadian oscillator. ZTL and ELF3 were reported to have opposite effects on clock function. *ztl* mutants and *ELF3* overexpression lines show a lengthened circadian period in light. Conversely *elf3* mutants and ZTL overexpression lines are reported as arrhythmic under constant light (LL) (Kim *et al.*, 2005). However, the *elf3-ztl* double mutant showed that ELF3 and ZTL have additive effects on the clock (Kim *et al.*, 2005). As GI controls the HSP90 mediated stabilisation of ZTL protein (Cha *et al.*, 2017, Kim *et al.*, 2011), ZTL protein then causes protein depletion of TOC1 via ubiquitination (Kim *et al.*, 2011). ELF3 interacts as a substrate adaptor for COP1 (an E3 ubiquitin ligase) to bind to and degrade GI protein, as a light input signal and indicator of day length in response to CRY2 (Yu *et al.*, 2008). The reduction of GI then prevents the formation of stable ZTL protein. Consequently this prevents ZTL-mediated inhibition of TOC1 in the central oscillator, and facilitates TOC1 action. ZTL also negatively regulates PRR5 by targeting PRR5 protein for degradation via the 26S proteasome (Fujiwara *et al.*, 2008, Kiba *et al.*, 2007). As PRR5 forms a negative regulatory feedback loop with LHY/ CCA1, ZTL therefore indirectly plays a role in the regulation of LHY/CCA1 within the central oscillator (Baudry *et al.*, 2010).

Far-red light

phyA is the presumed photoreceptor for detecting monochromatic far-red light. Mutations in *PHYA* resulted in loss of capacity for clock function (Wenden *et al.*, 2011). ELF4 was proposed to restrict far-red perception in those studies. Interestingly, the active form of phyA (phyA-Pfr) is formed under far-red light (Clough & Vierstra, 1997), given that far red converts the Pfr form of phytochrome back to the inactive Pr form. In part perhaps phyA evolved the ability to form Pfr under far red as a response to the change in light quality at the end of the day, which signals the transition from day to night and therefore the associated changes in environment. However, far red can also be a signal of shade due to far red being one of the only wavelengths of light able to pass through leaves (Federer & Tanner, 1966), suggesting that there may be different mechanisms to entrain the clock in these two different circumstances, as described in the next section. Plants in constant far-red light have a faster clock and show high expression of evening genes, such as *PRR1/TOC1*, and low expression of the morning genes *CCA1* and *LHY* (Wenden *et al.*, 2011). The exact mechanism of far-red input to the clock is not fully characterised. However, far red has been shown to be involved in the prevention of the interaction between Pfr and PIF3 (Martínez-García *et al.*, 2000). ELF4 was identified as playing a role in mediating far-red light input to the clock (Wenden *et al.*, 2011), Far-red light was used to aid recovery of rhythmicity in the otherwise arrhythmic *elf3* and *elf4* mutants (Kolmos *et al.*, 2011, Wenden *et al.*, 2011).

Shaded light

White light with supplementary far-red light causes the clock to slow down (Jiménez-Gómez *et al.*, 2010). Under shade, far red and potentially green light are present; there is a large overlap between far-red signalling and shade. Shade however, is a useful environmental indicator to plants for neighbour detection. *phyA* is thought to have the most involvement in mediating far-red signalling, but *phyB* also plays a key role (Kolmos *et al.*, 2011, Wenden *et al.*, 2011). Shading plants during the afternoon was found to have the greatest effect (Sellaro *et al.*, 2012). Responses to shade involve the degradation of phytochrome interacting factors, namely PIF4 and 5 (Lorrain *et al.*, 2008). PRR5 was found to regulate the shade-avoidance response by controlling PIF4 and PIF5, as well as downstream components of the phytochrome-mediated signalling pathway. Furthermore ZTL induces degradation of PRR5. However, this degradation was found to be repressed under blue light. It was suggested that PRR5 gates phytochrome mediated shade responses (Takase *et al.*, 2013). *ELF3* and *LUX* mutants (both components of the evening complex) show a reduced response to all wavelengths of light therefore growing with elongated hypocotyls as though under shade (Jiménez-Gómez *et al.*, 2010, Sellaro *et al.*, 2012, Zagotta *et al.*, 1996). This implies that *ELF3* and the evening-complex also play a role in the shade response to the clock (Kolmos *et al.*, 2011).

UV-B light

Ultraviolet B light (UVB) can be one of the more damaging wavelengths present in sunlight. UVB is a wavelength that is easily absorbed and damages both DNA and proteins (Jansen *et al.*, 1998), thus making UVB a useful light signal, but at the cost of inducing a stress response. UVB is an "anti-shade" signal informing a plant it is under direct sunlight. At lower-fluence rates, UV-B light is able to control development, promote photomorphogenesis, and drive gene expression (Heijde & Ulm, 2012). *Ultraviolet resistance locus 8 (UVR8)* drives signalling for the majority of UVB responses (Favory *et al.*, 2009, Rizzini *et al.*, 2011). Under UVB light, COP1 promotes the induction of elongated hypocotyl 5 (HY5) and HY5 homologue (HYH) which induce stress responses such as flavonoid biosynthesis to reduce UVB induced damage (Stracke *et al.*, 2010). UVR8 and COP1 are also crucial for UVB light entrainment of the clock (Fehér *et al.*, 2011). Under white light supplemented with UV-B light, COP1 induces HY5 and HYH, HY5 and HYH have not yet been implicated for clock entrainment by UVB (Fehér *et al.*, 2011). UVR8 is able to mediate both parametric and non-parametric entrainment, by inducing *PRR9* and *GI* under continuous light, alongside an increase in *CCA1* and *ELF3* response to UVB light pulses. UVR8 was identified as the UVB receptor that can mediate signal input to the oscillator, due to the fact that *uvr8* plants cannot input UVB light into the oscillator, (Fehér *et al.*, 2011, Heijde & Ulm, 2012). It has been proposed that UVR8 mediates UVB light input into the central oscillator by inhibiting PIF4 in the presence of UVB light. This requires COP1-mediated repression of *PIF4* transcript, and also through the stabilisation of HFR, which inhibits PIF4 (Hayes *et al.*, 2017). Canonical pathways used in UVB signalling mediate entrainment in the clock, but the critical nodes in entrainment are not fully resolved (Hayes *et al.*, 2017).

Green light

Many studies have been carried out to test the physiological effects that occur as a consequence of increased or absent green-light wavelengths. The mechanisms of sensing and input to the circadian clock are yet to be understood. It is thought that green wavelengths can operate via both a cryptochrome dependant and independent pathway (Folta & Maruhnich, 2007). Green light can reverse the effect of blue light on hypocotyl elongation (Bouly *et al.*, 2007, Folta, 2004), potentially due to the reversal of the blue light degradation of CRY1 (Bouly *et al.*, 2007). This could then have an effect on photoperiod and subsequently flowering time (Banerjee *et al.*, 2007, Folta & Maruhnich, 2007). The association of green light to cryptochromes was also shown by (Lin *et al.*, 1996) as overexpression of CRY1 causes increased sensitivity to green light. A reversible interaction between CRY and green light similar to that found for phytochrome in red and far red light suggests

that there are intermediate signalling factors similar to PIFs that are yet to be identified. It is thus plausible that green light could entrain the clock, but no definitive experiments have tested this.

Transcriptional regulation of photoreceptors by the clock.

The circadian clock generates rhythms of RNA and/or accumulation for all photoreceptor classes. Starting with the discovery that *phyB* mRNA is rhythmic (Bognár *et al.*, 1999, Toth *et al.*, 2001, Tóth *et al.*, 2001), subsequent findings revealed that all five phytochromes in Arabidopsis cycle. Interestingly, sub-nuclear accumulation of phytochrome holoprotein also appears to be under clock control. However, the implications of this are currently unclear. Similarly, *CRY* genes are rhythmic (Toth *et al.*, 2001). For *UVR8*, as UV light induces dimer disassembly to a monomer state, a diel cycle of dimers at night and monomers during the day occur (Findlay & Jenkins, 2016). *UVR8* mRNA displays robust circadian rhythms with a peak around subjective dusk (Mockler *et al.*, 2007). For *ZTL*, the mRNA generated does not cycle, but robust protein cycling is readily detectable. Together it is clear light receptors that act as input components to the clock, are themselves circadian-output regulated.

Phytochrome input to the central oscillator

Light input to the central oscillator is a daily *zeitgeber*, but the central oscillator also acts as a feedback mechanism to phytochromes over the day. The oscillator receives a number of light queues during the light phase of each day from photoreceptors detecting the different ratio of light wavelengths across the day. In turn, the oscillator inhibits expression of phytochrome proteins at points hypothesised in Figure 3.

At dawn (ZT0), *PHYC* peaks with the return of light and changing temperature (Toth *et al.*, 2001). HFR increases due to increased stability (Yang *et al.*, 2005). PIF4/5 expression peaks at dawn (Nomoto *et al.*, 2012), due to the lack of phytochromes, thus preventing phytochrome-induced degradation at dawn (Shin *et al.*, 2013). PIF4/5 continue to be present throughout the light phase, but are slowly degraded by interactions with the Pr form of phytochrome, PIFs can also interact with LHY, CCA1. PfrB interacts with PIF3, where PIF3 then binds to the G-box domain of *CCA1/LHY* promoters. CCA1 and LHY, as MYB transcription factors, then control other genes within the central oscillator, such as *PRR5*, 7 and 9, as well as non-circadian genes (Martínez-García *et al.*, 2000, Wang & Tobin, 1998).

PHYD and *E* are expressed two hours after dawn (Toth *et al.*, 2001). During the first half of the light phase (ZT0-6) there is an increase in light intensity up to ZT6, and alongside this, an increasing expression of phytochromes. *PHYB* and *CRY1* reach peaks in expression around mid-day (ZT6) as both *phyB* and *CRY1* work at high-light intensities (Lin *et al.*, 1996, Lin *et al.*, 1998, Toth *et al.*, 2001). *CAB1* expression peaks around mid-day as does the expression of HFR (Yang *et al.*, 2005). HFR is thought to interact with PIF3 (Fairchild *et al.*, 2000), but the mechanism through which this happens is not fully understood.

Both *PHYA* and *CRY2* peak towards the latter half of the light phase (ZT6-12) with the decreasing light and increasing far red intensity (Toth *et al.*, 2001). *CRY2* detects lower intensity light (Lin *et al.*, 1998), and the Pfr form of phytochrome is needed for *CRY2* activation (Mas *et al.*, 2000). *CRY2* then suppresses *PHYB* expression (Mas *et al.*, 2000). *phyA* is essential in controlling the clock in low light (Quail *et al.*, 1995, Somers *et al.*, 1998a) and far red conditions (Wenden *et al.*, 2011), potentially through the Pfr form of phytochrome being unable to interact with PIF3 (Martínez-García *et al.*, 2000).

At dusk, *CAB2* expression decreases, and *COP1* expression increases (Yang *et al.*, 2005), allowing *COP1* suppression of HFR throughout the dark phase of the day. *COP1* accumulation along with *ELF3* inhibits *GI* late in the afternoon (Yu *et al.*, 2008). *PILs* are rapidly produced during the first hour of shade, early into the dark phase and work with *TOC1* to restrict growth (Salter *et al.*, 2003). *PIF3* is at its highest level at dusk due to its interactions with the Pfr form of phytochrome and the highest level of Pfr being present just before dusk. Allowing information on high levels of far red light

to be input to the central oscillator (Martínez-García *et al.*, 2000). The evening complex (ELF3, ELF4, LUX) inhibits the transcription of *PIF4/5* at dusk (Herrero *et al.*, 2012, Nusinow *et al.*, 2011; Raschke *et al.* 2015). This allows PIF protein to accumulate stably due to the lack of phytochrome inhibition overnight, which thus promotes growth, reaching a maximal level at dawn (Shin *et al.*, 2013, Delker *et al.* 2014; Raschke *et al.* 2015).

COP1

Constitutive photomorphogenic 1 (COP1) is an E3 ubiquitin ligase, mediating day length input to the clock and flowering time. COP1 is negatively regulated by a direct protein-protein interaction with CRYs (Jang *et al.*, 2008, Wang *et al.*, 2001, Yang *et al.*, 2000). phyA and B affect the nuclear abundance of COP1 (Osterlund *et al.*, 1999), as the C-terminal domain of phyB directly interacts with COP1 (Millar *et al.*, 1994). COP1 acts as an intermediate, inputting photoperiodic information from PHY and CRY into the oscillator. COP1 in turn plays a negative regulatory role targeting phyA, phyB and HFR1 for ubiquitination (Osterlund *et al.*, 2000, Seo *et al.*, 2003, Yang *et al.*, 2005).

CRY1, CRY2, phyA and phyB all interact with COP1 via Suppressor of Phytochrome A (SPA). SPA1 is a nuclear localised repressor of phytochrome signalling (Hoecker *et al.*, 1999), which interacts with COP1 (Hoecker & Quail, 2001). SPA1 contains a coiled-coil domain that enhances the E3 ligase activity of COP1 on its targets (Seo *et al.*, 2003). The interactions between the 4 known SPA proteins and COP1, negatively regulate light signalling in response to certain wavelengths of light (Laubinger *et al.*, 2004, Zhu *et al.*, 2008). COP1-SPA1 interaction is repressed by activated CRY1 in blue light (Lian *et al.*, 2011), CRY2 interacts with COP1 via SPA1 to allow COP1 proteolysis of CO to control flowering time under blue light (Zuo *et al.*, 2011). The CRY1-SPA1 interaction enhances CRY2-SPA1 activity in response to blue light to suppress COP1 activity resulting in a suppression of CO degradation (Ordoñez-Herrera *et al.*, 2015, Zuo *et al.*, 2011).

In seedlings, phyA binds to SPA1 and 2 whereas, in adult plants phyA binds to SPA3 and 4 (Laubinger *et al.*, 2004). Binding of phyB to SPA1 is Pfr dependant allowing degradation of COP1/SPA1 in light conditions that promote nuclear accumulation of phyA and B, this enhances light responses, as the disruption of COP1/SPA1 interaction prevents degradation of photomorphogenic factors such as HFR, and HY5 (Sheerin *et al.*, 2015). The COP1/SPA complex is an important factor in repression of light responses in darkness, as the COP1/SPA complex interacts directly with photoreceptors leading to its inactivation (Huang *et al.*, 2014).

Within the central oscillator, COP1 interacts with ELF3 to mediate COP1 degradation of GI late in the afternoon (Yu *et al.*, 2008), potentially using ELF3 as an adaptor for COP1 binding to GI (Liu *et al.*, 2008). It is also possible that COP1 regulates the level of ELF3 present, in *cop1* mutants ELF3 protein accumulates to higher levels than in the wild type, but the mRNA levels remain unchanged (Liu *et al.*, 2001).

COP1 is also involved in UVB signalling as *cop1* mutants are deficient in a UVB response (Oravecz *et al.*, 2006). In the early stage of UVB signalling, UVR8 and COP1 directly interact in the nucleus (Favory *et al.*, 2009), UVR8 and COP1 were found to be essential for UVB entrainment (Fehér *et al.*, 2011). HY5 and HYH which are also important components of UVB signalling are regulated by COP1 (Brown & Jenkins, 2008). In the light COP1 detaches from HY5 allowing stabilisation and the light responsive target genes of HY5 to be activated (Yi & Deng, 2005). COP1 plays an important mediator role in the input of light from photoreceptors to the oscillator. In turn its regulation is dependent on photoreceptors, the short-period phenotype in mutant lines shows that COP1 plays a negative regulatory role on the clock.

PIFs and PILs

Phytochrome interacting factors (PIF) are a family of basic helix loop helix transcription factors. There are 4 well characterised PIFs, PIF1,3,4,5 (Leivar *et al.*, 2012). (Pfeiffer *et al.*, 2012) PIFs are unstable in the light due to their interaction with active phytochrome causing phosphorylation

and subsequent degradation (Leivar *et al.*, 2012, Soy *et al.*, 2012). The most well characterised PIF is PIF3, which was found to interact with the Pfr form of phytochrome B (PfrB) acting as a bridge between PfrB and its target gene by translocating PfrB to the nucleus. Thus allowing light induced control of gene expression, as PIF3 does not interact with the Pr form of phytochrome (Martínez-García *et al.*, 2000, Pfeiffer *et al.*, 2012). PIFs are also able to input information to the clock via direct interaction with clock genes that contain a G-box motif in their promoter, PIFs can interact directly with *LHY*, *CCA1*, *PRR5*, *PRR7*, *PRR9* and *LUX* (Martínez-García *et al.*, 2000). This is potentially one of the main mechanisms through which light/ day length information is used to control or alter the clock. The central oscillator in turn regulates PIF expression. Postdusk, *TOC1* peaks in expression, allowing direct interactions between *TOC1* and PIF3, which results in the *TOC1* gating of PIF induced growth, until *TOC1* levels decrease predawn (Soy *et al.*, 2016).

PIF4 and *5* show rhythmic expression with a diurnal peak at dawn (Nomoto *et al.*, 2012). Expression of *PIF4* and *5* is controlled by the evening complex, comprising of *ELF3*, *ELF4* and *LUX* (Herrero *et al.*, 2012), which binds to the promoter region of *PIF4* and *5* to inhibit transcription at dusk (Nusinow *et al.*, 2011). PIF protein stably accumulates overnight due to the lack of phytochrome induced degradation to reach their maximum level at dawn (Shin *et al.*, 2013). As PIFs are growth-promoting factors (Shin *et al.*, 2013), this leads to the highest growth rate occurring at the end of the night phase.

PIFs may also input information from other environmental cues to the clock such as temperature (McClung & Davis, 2010; Raschke *et al.* 2015). It was shown that *PIF4* expression also increases in response to temperature increases (Shin *et al.*, 2013). As dawn induces a temperature increase, alongside the return of daylight it would perhaps be advantageous to a plant to be able to input both of these environmental cues into the clock at the same time.

PIF3 like (PILs) are also basic helix loop helix transcription factors with large overlaps in function to PIFs, but have been associated with shade avoidance (Li *et al.*, 2014). This overlap in function has led to some ambiguous nomenclature as PIL5 is also referred to as PIF1 and likewise PIL6 as PIF5 (Li *et al.*, 2014). PIL1 has a distinct function and was shown to work with *TOC1* to restrict growth at specific times of day (Salter *et al.*, 2003). PIL1 accumulates rapidly within the first hour of shade cover acting as part of a rapid signalling pathway to stop growth (Li *et al.*, 2014), a secondary longer lasting shade response is then mediated by HFR and phytochrome rapidly regulated (*PAR1/2*) (Galstyan *et al.*, 2011). The exact mechanism through which PIL1 halts growth in shade is not known but a number of hypothesis were presented in (Li *et al.*, 2014). It was suggested that as PIF1 has a binding site for phyB it is possible that in shade PIL1 may outcompete PIF for DNA binding sites on the Pfr form of phyB, therefore reducing the growth promoting function of PIF5 (Li *et al.*, 2014). Alternatively, PIL1 may work via a PIF independent mechanism on components of downstream pathways; however this is yet to be tested.

HFR

Long hypocotyl in far red 1 (HFR1) is a basic helix loop helix transcription factor involved in phytochrome-mediated signalling (Fairchild *et al.*, 2000), and photomorphogenesis (Yang *et al.*, 2005). HFR is unstable in darkness and accumulates in the light, this accumulation is due to light preventing COP1 mediated degradation of HFR (Yang *et al.*, 2005). HFR is not able to bind phyA or B directly, instead HFR binds PIF3 forming potentially a heterodimer of PIF3/HFR which can then bind to the Pfr form of phyA/B. This is also highlighted by the fact that HFR is more abundantly found in far red light (Fairchild *et al.*, 2000). Mutants deficient in HFR had defective phyA responses, such as hypocotyl elongation, and induction of chlorophyll A binding protein (CAB) (Fankhauser & Chory, 2000). HFR is also thought to have a blue-light response (Duek & Fankhauser, 2003) through *CRY1* (Yang *et al.*, 2005), but the exact mechanism through which this occurs is not known.

Intersection of the clock components ELF3- and ELF4 to light and clock signalling: major integrators of light to the clock

EARLY FLOWERING 3 (ELF3) was first identified as a negative regulator of flowering time. In addition to the observation that *elf3* mutant was shown to be early flowering, large circadian defects were identified (Hicks *et al.*, 1996, Roden *et al.*, 2002; Undurraga *et al.*, 2012). *elf3* mutants are defective in gating of red-light perception to the clock. Cloning of ELF3 allowed for interactors to be detected, phyB was revealed to be a factor that associated to the N-terminus of ELF3 (Liu *et al.*, 2001). ELF3 was identified as playing a role in light signalling, in 12hours light 12 hours dark entrainment, ELF3 accumulates in the nucleus just before darkness (ZT12) (Liu *et al.*, 2001). Increasing day length, increases the nuclear accumulation of ELF3, increased darkness causes accumulation of ELF3 to decrease to an undetectable level (Liu *et al.*, 2001) showing a direct relationship between light and ELF3.

Phase response curves are made by measuring circadian period and phase during light pulses, at times across subjective day and night. Phase response curves for wild-type Arabidopsis in both red and blue light show the greatest effect and subsequent clock resetting to be caused by a light pulse during the subjective night. *ELF3* overexpression lines showed a much more gradual change in phase response with the same light pulses (Covington *et al.*, 2001). In white light, *ELF3* overexpression causes a period lengthening effect in a light-dependent manner. In darkness, the oscillator pace is not altered (Covington *et al.*, 2001). In *elf3* mutants, phase response light pulses showed a much greater effect than the wild type, suggesting that clock resetting is light dependent (Covington *et al.*, 2001), involving ELF3 in oscillator resetting by repressing the light input to the clock (Bujdoso & Davis, 2013). However, the exact mechanism through which this occurs is unknown.

elf3 and *phyB* mutants were found to have similar phenotypic traits, such as hypocotyl elongation in red light and constitutive shade avoidance (Devlin *et al.*, 1999). *elf3* mutants are also defective in their response to blue and/ or red light with stronger effects showing in red light. ELF3 grown in darkness is rhythmic with a long-period phenotype, rhythmic in light/dark entrainment cycles, but arrhythmic in free run light conditions (Hicks *et al.*, 1996). *elf3* plants are also unable to inhibit hypocotyl elongation under light (Zagotta *et al.*, 1996). The combined phenotypic characteristics of the *elf3* mutants suggest that ELF3 plays a role in light perception and signalling, particularly in red light due to the interaction between ELF3 and phyB (Reed *et al.*, 2000).

EARLY FLOWERING 4 (ELF4) was the first clock component interpreted as being required for the clock to cycle and it was revealed to be a component of normal light perception. Genetic loss of *ELF4* resulted in plants that were markedly attenuated in the ability of a red-light pulse to generate *CCA1* and *LHY* rhythms in etiolated plants. This was concluded to be due to ELF4 being required for the phytochrome-mediated light induction of *CCA1* and *LHY* expression (Kikis *et al.*, 2005). Consistent with this, it was shown that *elf4* mutants were hypermorphic and hypomorphic to red-light cues, dependent on the assay. Notably, red-light mediated induction of *CAB2* expression was elevated in *elf4* (McWatters *et al.*, 2007). This revealed that ELF4 contributes to so-called circadian gating of light responsiveness (negative photomorphogenesis), light-regulation of *PIF4/5* expression and the suppression of growth (positive photomorphogenesis) (Nozue *et al.*, 2007).

The presence of ELF4 in the nucleus increases the accumulation of ELF3 (Herrero *et al.*, 2012). ELF4 may function as a nuclear anchor for ELF3, but does not affect the nuclear localisation of LUX, the third component of the evening complex (Herrero *et al.*, 2012). Nuclear import of phyB is light dependent (Kircher *et al.*, 1999, Sakamoto & Nagatani, 1996), and it was shown that phyB does not import ELF3 into the nucleus (Bujdoso & Davis, 2013). It is possible that ELF3 plays a role in the nuclear import of phyB, as the N-terminus of ELF3 interacts with the C-terminal end of both the Pr and Pfr forms of PHYB (Liu *et al.*, 2001). However, ELF3 and phyB have opposite roles in controlling circadian oscillations (Herrero *et al.*, 2012). ELF3 needs the association with ELF4 to maintain circadian oscillations, counteracting the COP1 and phyB mediated repression of ELF3 (Herrero *et al.*, 2012). ELF3 is also part of the blue-light signalling pathway, through its interaction with COP1. How

phyB and ELF4 coordinate the action and localisation of ELF3 seems critical for the cooperative intersection of light perception and circadian clock function.

It was originally unclear what overall effect ELF3 has on other clock genes as the *elf3* loss of function mutation causes arrhythmicity. A reduction-of-function mutation in *elf3-12* provided a way to explore this, as the hypomorphic *elf3-12* allele is able to maintain rhythmicity (Kolmos *et al.*, 2011). This showed *elf3-12* to be light dependent but with a defective phase resetting mechanism. This *elf3-12* mutant allowed the position of ELF3 within the clock to be derived as key to the regulation of *PRR9* expression (Kolmos & Davis, 2007, Kolmos *et al.*, 2011). It is known that ELF3 associates to the promoter of *PRR9* to mediate its repression. In addition to clock-regulated *PRR9* transcription, *PRR9* expression is also light regulated. Furthermore, *prp9* mutants display photomorphogenic phenotypes (Nakamichi *et al.*, 2005). This highlights the role of ELF3 in reciprocally linking light signalling to clock function.

Overexpression of *PHYA* in an *elf3-12* background showed that the overexpressed *PHYA* has an additive effect with the *elf3-12* mutation to give a further shortened period in red light. In a range of light conditions the *elf3-12 PHYA*-overexpression lines had an altered phase however in darkness there was no change suggesting that light has an epistatic effect on *PHYA* (Kolmos *et al.*, 2011). Overexpression of *PHYB* in the same *elf3-12* background gave the same period shortening as the *PHYA*-overexpression lines; however the *PHYB-elf3-12* lines had a phase more closely linked to *PHYB* overexpression in a wild-type background (Kolmos *et al.*, 2011). This result suggests that *PHYB* functions upstream of *elf3-12* in light signalling and is able to suppress the effects of *elf3-12* (Bujdoso & Davis, 2013). Together it appears that some, but not all, inputs of phytochromes to the clock depend on ELF3.

Duration and quality of light

In regular light-dark cycles *Arabidopsis* has a circadian period of approximately 24 hours in light, whereas in darkness it has a period of 30 to 36 hours (Millar *et al.*, 1995). Light therefore makes the clock run faster, the absence of light cues causes the clock to slow, this is in keeping with Aschoff's rule (Aschoff, 1979). The range of photoreceptors present in *Arabidopsis* allow a range of fluence rates to be detected, ensuring the phase of the circadian oscillator is synchronised with environmental cues (Somers *et al.*, 1998a), at both dawn and dusk (Devlin & Kay, 2000a). Removal or even partial reduction of blue and red photoreceptors causes the clock to run slower (Millar *et al.*, 1995), suggesting that the effects of light intensity on the speed of the clock is limited by the number of photoreceptors present. This would also suggest that it would not be possible to increase the speed of the clock with higher light intensities beyond the maximum speed obtainable by that number of photoreceptors.

Input of duration and quality of light are important in synchronising processes such as flowering time and development (Weston *et al.*, 2000). Preceding photoperiod was shown to alter the subsequent speed of the clock (Darrah *et al.*, 2006; Boikoglou *et al.* 2011). Interestingly here is the long known role of ELF3 in processing light information to the clock (Hicks *et al.* 1996), and how this coordinates the capacity for a plant to perceive daily boundaries present in a day night cycle (McWatters *et al.* 2000). Recent work has revealed that extensive allelic variation at *ELF3* contribute to alterations in photoperiodic control and this is associated to alterations in encoded nuclear abundance and *in vivo* turn over diurnal time (Undurraga *et al.* 2012, Anwer *et al.* 2014). Combined with temperature variation over the day, light duration gives information on the time of year or season and therefore a warning of the growth conditions to follow. It is clear that allelic variation exists in *Arabidopsis* to change the output of such varying entrainment processes (Darrah *et al.*, 2006, Boikoglou *et al.* 2011; Anwer *et al.* 2014).

Plants are able to adapt to changes in light intensity, such as consistently low light intensity, by re-arranging photosynthetic machinery to be more efficient at light harvesting (Weston *et al.*, 2000). Blue light plays a major role in this. As such it would be assumed that cryptochromes and *PHYA/B* play a role in directing the timing of light capture. A connection hub for this, COP1 has been

identified as a signalling intermediate between these two processes (Walters *et al.*, 1999), however links between photoreceptor function and organisation of the photosynthetic apparatus await further investigation (Walters *et al.*, 1999, Weston *et al.*, 2000). Finally light intensity is also detected by the plastid sensing blue light, causing structural changes and elongation of the palisade to absorb more light, this process responds a lot more to blue light than red (Weston *et al.*, 2000). It could be hypothesised that this plastid information is used as a nuclear clock and is synchronised with the phyA and phyB red light input to the central oscillator by COP1.

Discussion

Multiple photoreceptors are essential components of light input to the clock. In this way they play a central role in the light input to the clock. Not only with the reversible, light-mediated reactions for maximal efficiency in light or dimmer light / shade, they also directly input light into the central oscillator through clock associated factors, such as ELF3/4, COP1, ZTL, PIFs, PILs and HFR. Transcriptional regulation and post-translational processes are all part of this complex web of interconnections between light-perception and clock function. Overall light input to the clock forms complex feedback systems that generated harmonised regulatory pathways, the mechanisms from light perception to clock function, and back again, awaits clear discoveries.

Many plant growth chambers are produced with red, blue and far-red LED panels, but are these the optimal light regimes to measure plant gene expression under? Furthermore, most chambers have a lights-on or off function which does not represent the graded changes in intensity that would occur with a plant growing under natural sunlight (or in a greenhouse), with the gradual appearance and disappearance of light at sunrise and sunset. As white light comprises a combination of different wavelengths of light simultaneously, it may be possible that the pathways for different colours of light interact more than has been found so far. Overlapping functions have been found between blue and red light and these are the most commonly used light wavelengths for plant LEDs. It is possible that there are essential components missing in just red and blue, although difficult to isolate in a complex web of circadian gene expression, may only be present in white light or in other light combinations not yet tested.

Light is essential for plant growth and it is therefore important to understand how plants process the daily light cues they receive. Further understanding of how each light wavelength is detected and the information fed into the central oscillator from each sensor, could potentially have a large impact on plant and ultimately crop growth. With global changes in climate, knowledge on essential lighting requirements and how this impacts on overall plant health could be used to optimise crop productivity. Indoor farming techniques using LEDs as a light source could be optimised for maximal yield and growth speed. As permafrost regions recede, and suitable growth land becomes available. A detailed understanding of photoperiodicity and how it impacts fitness will help with the challenges created by growing crops in shorter growth seasons and longer daylight hours.

Figure legends

Figure 1. Schematic diagram of feedback loops in the Arabidopsis central oscillator.

Figure 2. Effects of light intensity on circadian period. A). Circadian gene expression is rhythmic in constant light but in the absence of exogenous sucrose, rapidly becomes arrhythmic in prolonged darkness. B). Following Aschoff's rule, high intensity light causes the clock to run faster; lower light intensities cause the clock to slow. C). High light intensities result in period shortening; lower light intensities result in period lengthening.

Figure 3. Circadian photoreceptors and their potential interactions to input light to the central oscillator.

For Review Only

References

- Adams S., Manfield I., Stockley P. & Carré I.A. (2015) Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. *PLOS ONE*, **10**, e0143943.
- Ahmad M. & Cashmore A.R. (1993) HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature*, **366**, 162-166.
- Ahmad M., Jarillo J.A. & Cashmore A.R. (1998a) Chimeric Proteins between cry1 and cry2 Arabidopsis Blue Light Photoreceptors Indicate Overlapping Functions and Varying Protein Stability. *The Plant Cell*, **10**, 197-207.
- Ahmad M., Jarillo J.A., Smirnova O. & Cashmore A.R. (1998b) The CRY1 Blue Light Photoreceptor of Arabidopsis Interacts with Phytochrome A In Vitro. *Molecular Cell*, **1**, 939-948.
- Alabadí D., Oyama T., Yanovsky M.J., Harmon F.G., Más P. & Kay S.A. (2001) Reciprocal Regulation Between TOC1 and LHY/CCA1 Within the Arabidopsis Circadian Clock. *Science*, **293**, 880-883.
- Anderson S.L., Somers D.E., Millar A.J., Hanson K., Chory J. & Kay S.A. (1997) Attenuation of phytochrome A and B signaling pathways by the Arabidopsis circadian clock. *The Plant Cell*, **9**, 1727-1743.
- Anwer M.U., Boikoglou E., Herrero E., Hallstein M., Davis A.M., Velikkakam James G., Nagy F. & Davis S.J. (2014) Natural variation reveals that intracellular distribution of ELF3 protein is associated with function in the circadian clock. *eLife*, e02206.
- Aschoff J. (1979) Circadian Rhythms: Influences of Internal and External Factors on the Period Measured in Constant Conditions. *Zeitschrift für Tierpsychologie*, **49**, 225-249.
- Aukerman M.J., Hirschfeld M., Wester L., Weaver M., Clack T., Amasino R.M. & Sharrock R.A. (1997) A deletion in the PHYD gene of the Arabidopsis Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *The Plant Cell*, **9**, 1317-1326.
- Banerjee R., Schleicher E., Meier S., Viana R.M., Pokorný R., Ahmad M., Bittl R. & Batschauer A. (2007) The Signaling State of Arabidopsis Cryptochrome 2 Contains Flavin Semiquinone. *Journal of Biological Chemistry*, **282**, 14916-14922.
- Baudry A., Ito S., Song Y.H., Strait A.A., Kiba T., Lu S., Henriques R., Pruneda-Paz J.L., Chua N.H., Tobin E.M., Kay S.A., Imaizumi T. (2010) F-box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control Arabidopsis clock progression. *The Plant Cell*, **22**, 606-622.
- Bläsing O.E., Gibon Y., Günther M., Höhne M., Morcuende R., Osuna D., Thimm O., Usadel B., Scheible W.-R. & Stitt M. (2005) Sugars and Circadian Regulation Make Major Contributions to the Global Regulation of Diurnal Gene Expression in Arabidopsis. *The Plant Cell*, **17**, 3257-3281.
- Bognár L.K., Hall A., Ádám É., Thain S.C., Nagy F. & Millar A.J. (1999) The circadian clock controls the expression pattern of the circadian input photoreceptor, phytochrome B. *Proceedings of the National Academy of Sciences of the United States of America*, **96**, 14652-14657.
- Boikoglou E., Ma Z., von Korff M., Davis A.M., Nagy F. & Davis S.J. (2011) Environmental memory from a circadian oscillator: the Arabidopsis thaliana clock differentially integrates perception of photic vs. thermal entrainment. *Genetics*, **189**, 655-664.
- Bouly J.P., Schleicher E., Dionisio-Sese M., Vandenbussche F., Van Der Straeten D., Bakrim N., Meier S., Batschauer A., Galland P., Bittl R. & Ahmad M. (2007) Cryptochrome Blue Light Photoreceptors Are Activated through Interconversion of Flavin Redox States. *Journal of Biological Chemistry*, **282**, 9383-9391.
- Brown B.A. & Jenkins G.I. (2008) UV-B Signaling Pathways with Different Fluence-Rate Response Profiles Are Distinguished in Mature Arabidopsis Leaf Tissue by Requirement for UVR8, HY5, and HYH. *Plant Physiology*, **146**, 576-588.
- Bujdoso N. & Davis S. (2013) Mathematical modeling of an oscillating gene circuit to unravel the circadian clock network of Arabidopsis thaliana. *Frontiers in Plant Science*, **4**.

- 691 Carre I.A. & Kay S.A. (1995) Multiple DNA-Protein Complexes at a Circadian-Regulated Promoter
692 Element. *The Plant Cell*, **7**, 2039-2051.
- 693 Cashmore A.R., Jarillo J.A., Wu Y.J. & Liu D. (1999) Cryptochromes: Blue Light Receptors for Plants
694 and Animals. *Science*, **284**, 760-765.
- 695 Cha J.Y., Kim J., Kim T.S., Zeng Q., Wang L., Lee S.Y., Kim W.Y. & Somers D.E. (2017) GIGANTEA is a
696 co-chaperone which facilitates maturation of ZEITLUPE in the Arabidopsis circadian clock.
697 *Nature Communications*, **8**, 3.
- 698 Chow B.Y., Sanchez S.E., Breton G., Pruneda-Paz J.L., Krogan N.T. & Kay S.A. (2014) Transcriptional
699 Regulation of LUX by CBF1 Mediates Cold Input to the Circadian Clock in Arabidopsis. *Current*
700 *Biology*, **24**, 1518-1524.
- 701 Clough R.C. & Vierstra R.D. (1997) Phytochrome degradation. *Plant, Cell & Environment*, **20**, 713-721.
- 702 Covington M.F., Panda S., Liu X.L., Strayer C.A., Wagner D.R. & Kay S.A. (2001) ELF3 Modulates
703 Resetting of the Circadian Clock in Arabidopsis. *The Plant Cell*, **13**, 1305-1316.
- 704 Darrah C., Taylor B.L., Edwards K.D., Brown P.E., Hall A. & McWatters H.G. (2006) Analysis of Phase
705 of LUCIFERASE Expression Reveals Novel Circadian Quantitative Trait Loci in Arabidopsis.
706 *Plant Physiology*, **140**, 1464-1474.
- 707 Davis S.J. (2002) Photoperiodism: The Coincidental Perception of the Season. *Current Biology*, **12**,
708 R841-R843.
- 709 Delker C., Sonntag L., James G.V., Janitza P., Ibañez C., Ziermann H., Peterson T., Denk K., Mull S.,
710 Ziegler J., Davis S.J., Schneeberger K. & Quint M. (2014) The DET1-COP1-HY5 pathway
711 constitutes a multipurpose signaling module regulating plant photomorphogenesis and
712 thermomorphogenesis. *Cell Rep.* **24**:1983-1989.
- 713 Devlin P.F. & Kay S.A. (2000a) Cryptochromes are required for phytochrome signaling to the
714 circadian clock but not for rhythmicity. *Plant Cell*, **12**, 2499-2510.
- 715 Devlin P.F. & Kay S.A. (2000b) Flower arranging in Arabidopsis. *Science*, **288**, 1600-1602.
- 716 Devlin P.F., Patel S.R. & Whitelam G.C. (1998) Phytochrome E Influences Internode Elongation and
717 Flowering Time in Arabidopsis. *The Plant Cell*, **10**, 1479-1487.
- 718 Devlin P.F., Robson P.R.H., Patel S.R., Goosey L., Sharrock R.A. & Whitelam G.C. (1999) Phytochrome
719 D Acts in the Shade-Avoidance Syndrome in Arabidopsis by Controlling Elongation Growth
720 and Flowering Time. *Plant Physiology*, **119**, 909-916.
- 721 Dodd A.N., Salathia N., Hall A., Kevei E., Toth R., Nagy F., Hibberd J.M., Millar A.J. & Webb A.A.R.
722 (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive
723 advantage. *Science*, **309**, 630-633.
- 724 Duek P.D. & Fankhauser C. (2003) HFR1, a putative bHLH transcription factor, mediates both
725 phytochrome A and cryptochrome signalling. *The Plant Journal*, **34**, 827-836.
- 726 Edwards K.D., Guerineau F., Devlin P.F. & Millar A.J. (2015) Low temperature effects of
727 *PHYTOCHROME C* on the circadian clock in Arabidopsis suggest that *PHYC* underlies natural
728 variation in biological timing. *bioRxiv*.
- 729 Fairchild C.D., Schumaker M.A. & Quail P.H. (2000) HFR1 encodes an atypical bHLH protein that acts
730 in phytochrome A signal transduction. *Genes & Development*, **14**, 2377-2391.
- 731 Fankhauser C. & Chory J. (2000) RSF1, an Arabidopsis Locus Implicated in Phytochrome A Signaling.
732 *Plant Physiology*, **124**, 39-46.
- 733 Favory J.J., Stec A., Gruber H., Rizzini L., Oravec A., Funk M., Albert A., Cloix C., Jenkins G.I., Oakeley
734 E.J., Seidlitz H.K., Nagy F. & Ulm R. (2009) Interaction of COP1 and UVR8 regulates UV-B-
735 induced photomorphogenesis and stress acclimation in Arabidopsis. *The EMBO Journal*, **28**,
736 591-601.
- 737 Federer C.A. & Tanner C.B. (1966) Spectral Distribution of Light in the Forest. *Ecology*, **47**, 555-560.
- 738 Fehér B., Kozma-Bognár L., Kevei É., Hajdu A., Binkert M., Davis S.J., Schäfer E., Ulm R. & Nagy F.
739 (2011) Functional interaction of the circadian clock and UV RESISTANCE LOCUS 8-controlled
740 UV-B signaling pathways in Arabidopsis thaliana. *The Plant Journal*, **67**, 37-48.

- Findlay K.M.W. & Jenkins G.I. (2016) Regulation of UVR8 photoreceptor dimer/monomer photo-equilibrium in Arabidopsis plants grown under photoperiodic conditions. *Plant, Cell & Environment*, **39**, 1706-1714.
- Folta K.M. (2004) Green Light Stimulates Early Stem Elongation, Antagonizing Light-Mediated Growth Inhibition. *Plant Physiology*, **135**, 1407-1416.
- Folta K.M. & Maruhnich S.A. (2007) Green light: a signal to slow down or stop. *Journal of Experimental Botany*, **58**, 3099-3111.
- Fornara F., de Montaigu A., Sanchez-Villarreal A., Takahashi Y., van Themaat E.V.L., Huettel B., Davis S.J. & Coupland G. (2015) The GI-CDF module of Arabidopsis affects freezing tolerance and growth as well as flowering. *Plant Journal*, **81**, 695-706.
- Franklin K.A., Davis S.J., Stoddart W.M., Vierstra R.D. & Whitelam G.C. (2003) Mutant Analyses Define Multiple Roles for Phytochrome C in Arabidopsis Photomorphogenesis. *The Plant Cell*, **15**, 1981-1989.
- Franklin K.A. & Quail P.H. (2010) Phytochrome functions in Arabidopsis development. *Journal of Experimental Botany*, **61**, 11-24.
- Fujiwara S., Wang L., Han L., Suh S. S., Salomé P. A., McClung C. R., Somers D. E. (2008) Post-translational regulation of the Arabidopsis circadian clock through selective proteolysis and phosphorylation of pseudo-response regulator proteins. *Journal of Biological Chemistry*, **283**, 23073-23083.
- Galstyan A., Cifuentes-Esquivel N., Bou-Torrent J. & Martinez-Garcia J.F. (2011) The shade avoidance syndrome in Arabidopsis: a fundamental role for atypical basic helix-loop-helix proteins as transcriptional cofactors. *The Plant Journal*, **66**, 258-267.
- Gendron J.M., Pruneda-Paz J.L., Doherty C.J., Gross A.M., Kang S.E. & Kay S.A. (2012) Arabidopsis circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proceedings of the National Academy of Sciences*, **109**, 3167-3172.
- Goto N., Kumagai T. & Koornneef M. (1991) Flowering responses to light-breaks in photomorphogenic mutants of Arabidopsis thaliana, a long-day plant. *Physiologia Plantarum*, **83**, 209-215.
- Guerriero M.L., Pokhilko A., Fernández A.P., Halliday K.J., Millar A.J. & Hillston J. (2012) Stochastic properties of the plant circadian clock. *Journal of the Royal Society Interface*, **9**, 744-756.
- Guo H., Yang H., Mockler T.C. & Lin C. (1998) Regulation of Flowering Time by Arabidopsis Photoreceptors. *Science*, **279**, 1360-1363.
- Habte E., Muller L.M., Shtaya M., Davis S.J. & von Korff M. (2014) Osmotic stress at the barley root affects expression of circadian clock genes in the shoot. *Plant Cell and Environment*, **37**, 1321-1337.
- Hanano S., Stracke R., Jakoby M., Merkle T., Domagalska M.A., Weisshaar B. & Davis S.J. (2008) A systematic survey in Arabidopsis thaliana of transcription factors that modulate circadian parameters. *BMC Genomics*, **9**, 182.
- Harmer S.L. (2009) The Circadian System in Higher Plants. *Annual Review of Plant Biology*, **60**, 357-377.
- Harmer S.L., Hogenesch J.B., Straume M., Chang H.S., Han B., Zhu T., Wang X., Kreps J.A. & Kay S.A. (2000) Orchestrated Transcription of Key Pathways in Arabidopsis by the Circadian Clock. *Science*, **290**, 2110-2113.
- Haydon M.J., Mielczarek O., Robertson F.C., Hubbard K.E. & Webb A.A.R. (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. *Nature*, **502**, 689-692.
- Haydon M.J. & Webb A.A. (2016) Assessing the Impact of Photosynthetic Sugars on the Arabidopsis Circadian Clock. *Environmental Responses in Plants: Methods and Protocols*, 133-140.
- Hayes S., Sharma A., Fraser D.P., Trevisan M., Cragg-Barber C.K., Tavridou E., Fankhauser C., Jenkins G.I. & Franklin K.A. (2017) UV-B Perceived by the UVR8 Photoreceptor Inhibits Plant Thermomorphogenesis. *Current Biology*, **27**, 120-127.

- 791 Hazen S.P., Schultz T.F., Pruneda-Paz J.L., Borevitz J.O., Ecker J.R. & Kay S.A. (2005) LUX ARRHYTHMO
792 encodes a Myb domain protein essential for circadian rhythms. *Proceedings of the National*
793 *Academy of Sciences of the United States of America*, **102**, 10387-10392.
- 794 Heijde M. & Ulm R. (2012) UV-B photoreceptor-mediated signalling in plants. *Trends in Plant Science*,
795 **17**, 230-237.
- 796 Helfer A., Nusinow D.A., Chow B.Y., Gehrke A.R., Bulyk M.L. & Kay S.A. (2011) LUX ARRHYTHMO
797 Encodes a Nighttime Repressor of Circadian Gene Expression in the Arabidopsis Core Clock.
798 *Current Biology*, **21**, 126-133.
- 799 Herrero E., Kolmos E., Bujdosó N., Yuan Y., Wang M., Berns M.C., Uhlworm H., Coupland G., Saini R.,
800 Jaskolski M., Webb A., Gonçalves J. & Davis S.J. (2012) EARLY FLOWERING4 Recruitment of
801 EARLY FLOWERING3 in the Nucleus Sustains the Arabidopsis Circadian Clock. *The Plant Cell*,
802 **24**, 428-443.
- 803 Hicks K.A., Millar A.J., Carré I.A., Somers D.E., Straume M., Meeks-Wagner D.R. & Kay S.A. (1996)
804 Conditional Circadian Dysfunction of the Arabidopsis early-flowering 3 Mutant. *Science*, **274**,
805 790-792.
- 806 Hoecker U. & Quail P.H. (2001) The Phytochrome A-specific Signaling Intermediate SPA1 Interacts
807 Directly with COP1, a Constitutive Repressor of Light Signaling in Arabidopsis. *Journal of*
808 *Biological Chemistry*, **276**, 38173-38178.
- 809 Hoecker U., Tepperman J.M. & Quail P.H. (1999) SPA1, a WD-Repeat Protein Specific to
810 Phytochrome A Signal Transduction. *Science*, **284**, 496.
- 811 Hu W., Franklin K.A., Sharrock R.A., Jones M.A., Harmer S.L. & Lagarias J.C. (2013) Unanticipated
812 regulatory roles for Arabidopsis phytochromes revealed by null mutant analysis. *Proceedings*
813 *of the National Academy of Sciences*, **110**, 1542-1547.
- 814 Huang H., Alvarez S., Bindbeutel R., Shen Z., Naldrett M.J., Evans B.S., Briggs S.P., Hicks L.M., Kay S.A.
815 & Nusinow D.A. (2016) Identification of Evening Complex Associated Proteins in Arabidopsis
816 by Affinity Purification and Mass Spectrometry. *Molecular & Cellular Proteomics*, **15**, 201-
817 217.
- 818 Huang X., Ouyang X. & Deng X.W. (2014) Beyond repression of photomorphogenesis: role switching
819 of COP/DET/FUS in light signaling. *Current Opinion in Plant Biology*, **21**, 96-103.
- 820 Jang S., Marchal V., Panigrahi K.C.S., Wenkel S., Soppe W., Deng X.-W., Valverde F. & Coupland G.
821 (2008) Arabidopsis COP1 shapes the temporal pattern of CO accumulation conferring a
822 photoperiodic flowering response. *The EMBO Journal*, **27**, 1277-1288.
- 823 Jansen M.A.K., Gaba V. & Greenberg B.M. (1998) Higher plants and UV-B radiation: balancing
824 damage, repair and acclimation. *Trends in Plant Science*, **3**, 131-135.
- 825 Jarillo J.A., Capel J., Tang R.H., Yang H.Q., Alonso J.M., Ecker J.R. & Cashmore A.R. (2001) An
826 Arabidopsis circadian clock component interacts with both CRY1 and phyB. *Nature*, **410**, 487-
827 490.
- 828 Jiménez-Gómez J.M., Wallace A.D. & Maloof J.N. (2010) Network analysis identifies ELF3 as a QTL for
829 the shade avoidance response in Arabidopsis. *PLoS Genet*, **6**, e1001100.
- 830 Johnson C.H., Knight M.R., Kondo T., Masson P., Sedbrook J., Haley A. & Trewavas A. (1995) Circadian
831 oscillations of cytosolic and chloroplastic free calcium in plants. *Science*, **269**, 1863.
- 832 Kamioka M., Takao S., Suzuki T., Taki K., Higashiyama T., Kinoshita T. & Nakamichi N. (2016) Direct
833 Repression of Evening Genes by CIRCADIANT CLOCK-ASSOCIATED1 in the Arabidopsis
834 Circadian Clock. *The Plant Cell*, **28**, 696-711.
- 835 Kevei É., Gyula P., Hall A., Kozma-Bognár L., Kim W.Y., Eriksson M.E., Tóth R., Hanano S., Fehér B.,
836 Southern M.M., Bastow R.M., Viczián A., Hibberd V., Davis S.J., Somers D.E., Nagy F. & Millar
837 A.J. (2006) Forward Genetic Analysis of the Circadian Clock Separates the Multiple Functions
838 of ZEITLUPE. *Plant Physiology*, **140**, 933-945.
- 839 Kiba T., Henriques R., Hitoshi S., Chua, N.H. (2007) Targeted Degradation of PSEUDO-RESPONSE
840 REGULATOR5 by an SCF^{ZTL} Complex Regulates Clock Function and Photomorphogenesis in
841 Arabidopsis thaliana. *The Plant Cell*, **19** (8) 2516-2530.

- 842 Kikis E.A., Khanna R. & Quail P.H. (2005) ELF4 is a phytochrome-regulated component of a negative-
 843 feedback loop involving the central oscillator components CCA1 and LHY. *The Plant Journal*,
 844 **44**, 300-313.
- 845 Kim T.s., Kim W.Y., Fujiwara S., Kim J., Cha J.Y., Park J.H., Lee S.Y. & Somers D.E. (2011) HSP90
 846 functions in the circadian clock through stabilization of the client F-box protein ZEITLUPE.
 847 *Proceedings of the National Academy of Sciences*, **108**, 16843-16848.
- 848 Kim W.Y., Fujiwara S., Suh S.S., Kim J., Kim Y., Han L., David K., Putterill J., Nam H.G. & Somers D.E.
 849 (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature*,
 850 **449**, 356-360.
- 851 Kim W.Y., Hicks K.A. & Somers D.E. (2005) Independent Roles for EARLY FLOWERING 3 and ZEITLUPE
 852 in the Control of Circadian Timing, Hypocotyl Length, and Flowering Time. *Plant Physiology*,
 853 **139**, 1557-1569.
- 854 Kircher S., Kozma-Bognár L., Kim L., Adam E., Harter K., Schäfer E. & Nagy F. (1999) Light Quality-
 855 Dependent Nuclear Import of the Plant Photoreceptors Phytochrome A and B. *The Plant Cell*,
 856 **11**, 1445-1456.
- 857 Kolmos E. & Davis S.J. (2007) ELF4 as a Central Gene in the Circadian Clock. *Plant Signaling &*
 858 *Behavior*, **2**, 370-372.
- 859 Kolmos E., Herrero E., Bujdosó N., Millar A.J., Toth R., Gyula P., Nagy F. & Davis S.J. (2011) A
 860 Reduced-Function Allele Reveals That EARLY FLOWERING3 Repressive Action on the
 861 Circadian Clock Is Modulated by Phytochrome Signals in Arabidopsis. *The Plant Cell*, **23**,
 862 3230-3246.
- 863 Koornneef M., Rolff E. & Spruit C.J.P. (1980) Genetic Control of Light-inhibited Hypocotyl Elongation
 864 in Arabidopsis thaliana (L.) Heynh. *Zeitschrift für Pflanzenphysiologie*, **100**, 147-160.
- 865 Laubinger S., Fittinghoff K. & Hoecker U. (2004) The SPA Quartet: A Family of WD-Repeat Proteins
 866 with a Central Role in Suppression of Photomorphogenesis in Arabidopsis. *The Plant Cell*, **16**,
 867 2293-2306.
- 868 Leivar P., Monte E., Cohn M.M. & Quail P.H. (2012) Phytochrome Signaling in Green Arabidopsis
 869 Seedlings: Impact Assessment of a Mutually Negative phyB–PIF Feedback Loop. *Molecular*
 870 *Plant*, **5**, 734-749.
- 871 Li L., Zhang Q., Pedmale U.V., Nito K., Fu W., Lin L., Hazen S.P. & Chory J. (2014) PIL1 Participates in a
 872 Negative Feedback Loop that Regulates Its Own Gene Expression in Response to Shade.
 873 *Molecular Plant*, **7**, 1582-1585.
- 874 Lian H.L., He S.B., Zhang Y.C., Zhu D.M., Zhang J.Y., Jia K.P., Sun S.X., Li L. & Yang H.Q. (2011) Blue-
 875 light-dependent interaction of cryptochrome 1 with SPA1 defines a dynamic signaling
 876 mechanism. *Genes & Development*, **25**, 1023-1028.
- 877 Lin C., Ahmad M. & Cashmore A.R. (1996) Arabidopsis cryptochrome 1 is a soluble protein mediating
 878 blue light-dependent regulation of plant growth and development. *The Plant Journal*, **10**,
 879 893-902.
- 880 Lin C., Yang H., Guo H., Mockler T., Chen J. & Cashmore A.R. (1998) Enhancement of blue-light
 881 sensitivity of Arabidopsis seedlings by a blue light receptor cryptochrome 2. *Proceedings of*
 882 *the National Academy of Sciences*, **95**, 2686-2690.
- 883 Liu H., Yu X., Li K., Klejnot J., Yang H., Lisiero D. & Lin C. (2008) Photoexcited CRY2 Interacts with CIB1
 884 to Regulate Transcription and Floral Initiation in Arabidopsis. *Science*, **322**, 1535-1539.
- 885 Liu X.L., Covington M.F., Fankhauser C., Chory J. & Wagner D.R. (2001) ELF3 Encodes a Circadian
 886 Clock–Regulated Nuclear Protein That Functions in an Arabidopsis PHYB Signal Transduction
 887 Pathway. *The Plant Cell*, **13**, 1293-1304.
- 888 Locke J.C.W., Kozma-Bognár L., Gould P.D., Fehér B., Kevei É., Nagy F., Turner M.S., Hall A. & Millar
 889 A.J. (2006) Experimental validation of a predicted feedback loop in the multi-oscillator clock
 890 of Arabidopsis thaliana. *Molecular Systems Biology*, **2**.


- 891 Lorrain S., Allen T., Duek P.D., Whitelam G.C. & Fankhauser C. (2008) Phytochrome-mediated
892 inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription
893 factors. *The Plant Journal*, **53**, 312-323.
- 894 Martínez-García J.F., Huq E. & Quail P.H. (2000) Direct Targeting of Light Signals to a Promoter
895 Element-Bound Transcription Factor. *Science*, **288**, 859-863.
- 896 Más P., Alabadí D., Yanovsky M.J., Oyama T. & Kay S.A. (2003) Dual Role of TOC1 in the Control of
897 Circadian and Photomorphogenic Responses in Arabidopsis. *The Plant Cell*, **15**, 223-236.
- 898 Mas P., Devlin P.F., Panda S. & Kay S.A. (2000) Functional interaction of phytochrome B and
899 cryptochrome 2. *Nature*, **408**, 207-211.
- 900 Mas P., Kim W.Y., Somers D.E. & Kay S.A. (2003) Targeted degradation of TOC1 by ZTL modulates
901 circadian function in Arabidopsis thaliana. *Nature*, **426**, 567-570.
- 902 Mathews S. & Sharrock R.A. (1997) Phytochrome gene diversity. *Plant, Cell & Environment*, **20**, 666-
903 671.
- 904 McClung C.R. & Davis S.J. (2010) Ambient Thermometers in Plants: From Physiological Outputs
905 towards Mechanisms of Thermal Sensing. *Current Biology*, **20**, R1086-R1092.
- 906 McWatters H.G., Bastow R.M., Hall A. & Millar A.J. (2000) The ELF3 zeitnehmer regulates light
907 signalling to the circadian clock. *Nature*, **408**:716-720.
- 908 McWatters H.G., Kolmos E., Hall A., Doyle M.R., Amasino R.M., Gyula P., Nagy F., Millar A.J. & Davis
909 S.J. (2007) ELF4 Is Required for Oscillatory Properties of the Circadian Clock. *Plant*
910 *Physiology*, **144**, 391-401.
- 911 Michael T.P., Salomé P.A., Yu H.J., Spencer T.R., Sharp E.L., McPeck M.A., Alonso J.M., Ecker J.R. &
912 McClung C.R. (2003) Enhanced Fitness Conferred by Naturally Occurring Variation in the
913 Circadian Clock. *Science*, **302**, 1049-1053.
- 914 Millar A., Straume M., Chory J., Chua N. & Kay S. (1995) The regulation of circadian period by
915 phototransduction pathways in Arabidopsis. *Science*, **267**, 1163-1166.
- 916 Millar A.J. & Kay S.A. (1996) Integration of circadian and phototransduction pathways in the network
917 controlling CAB gene transcription in Arabidopsis. *Proceedings of the National Academy of*
918 *Sciences*, **93**, 15491-15496.
- 919 Millar A.J., McGrath R.B. & Chua N.H. (1994) PHYTOCHROME PHOTOTRANSDUCTION PATHWAYS.
920 *Annual Review of Genetics*, **28**, 325-349.
- 921 Mizoguchi T., Wright L., Fujiwara S., Cremer F., Lee K., Onouchi H., Mouradov A., Fowler S., Kamada
922 H., Putterill J. & Coupland G. (2005) Distinct Roles of GIGANTEA in Promoting Flowering and
923 Regulating Circadian Rhythms in Arabidopsis. *The Plant Cell*, **17**, 2255-2270.
- 924 Mizuno T., Nomoto Y., Oka H., Kitayama M., Takeuchi A., Tsubouchi M. & Yamashino T. (2014)
925 Ambient temperature signal feeds into the circadian clock transcriptional circuitry through
926 the EC night-time repressor in Arabidopsis thaliana. *Plant and Cell Physiology*, **55**, 958-976.
- 927 Mockler T., Michael T., Priest H., Shen R., Sullivan C., Givan S., McEntee C., Kay S. & Chory J. (2007)
928 *The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern*
929 *matching, and promoter analysis*. Paper presented at the Cold Spring Harbor Symposia on
930 Quantitative Biology.
- 931 Nakamichi N., Kita M., Ito S., Yamashino T. & Mizuno T. (2005) PSEUDO-RESPONSE REGULATORS,
932 PRR9, PRR7 and PRR5, Together Play Essential Roles Close to the Circadian Clock of
933 Arabidopsis thaliana. *Plant and Cell Physiology*, **46**, 686-698.
- 934 Nakamichi N., Kusano M., Fukushima A., Kita M., Ito S., Yamashino T., Saito K., Sakakibara H. &
935 Mizuno T. (2009) Transcript Profiling of an Arabidopsis PSEUDO RESPONSE REGULATOR
936 Arrhythmic Triple Mutant Reveals a Role for the Circadian Clock in Cold Stress Response.
937 *Plant and Cell Physiology*, **50**, 447-462.
- 938 Nomoto Y., Kubozono S., Yamashino T., Nakamichi N. & Mizuno T. (2012) Circadian Clock- and PIF4-
939 Controlled Plant Growth: A Coincidence Mechanism Directly Integrates a Hormone Signaling
940 Network into the Photoperiodic Control of Plant Architectures in Arabidopsis thaliana. *Plant*
941 *and Cell Physiology*, **53**, 1950-1964.


- 942 Nozue K., Covington M.F., Duek P.D., Lorrain S., Fankhauser C., Harmer S.L. & Maloof J.N. (2007)
943 Rhythmic growth explained by coincidence between internal and external cues. *Nature*, **448**,
944 358-361.
- 945 Nusinow D.A., Helfer A., Hamilton E.E., King J.J., Imaizumi T., Schultz T.F., Farre E.M. & Kay S.A.
946 (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl
947 growth. *Nature*, **475**, 398-402.
- 948 Onai K. & Ishiura M. (2005) PHYTOCLOCK 1 encoding a novel GARP protein essential for the
949 Arabidopsis circadian clock. *Genes to Cells*, **10**, 963-972.
- 950 Oravecz A., Baumann A., Máté Z., Brzezinska A., Molinier J., Oakeley E.J., Ádám É., Schäfer E., Nagy F.
951 & Ulm R. (2006) CONSTITUTIVELY PHOTOMORPHOGENIC1 Is Required for the UV-B
952 Response in Arabidopsis. *The Plant Cell*, **18**, 1975-1990.
- 953 Ordoñez-Herrera N., Fackendahl P., Yu X., Schaefer S., Koncz C. & Hoecker U. (2015) A cop1 spa
954 Mutant Deficient in COP1 and SPA Proteins Reveals Partial Co-Action of COP1 and SPA
955 during Arabidopsis Post-Embryonic Development and Photomorphogenesis. *Molecular*
956 *Plant*, **8**, 479-481.
- 957 Osterlund M.T., Ang L.H. & Deng X.W. (1999) The role of COP1 in repression of Arabidopsis
958 photomorphogenic development. *Trends in Cell Biology*, **9**, 113-118.
- 959 Osterlund M.T., Hardtke C.S., Wei N. & Deng X.W. (2000) Targeted destabilization of HY5 during
960 light-regulated development of Arabidopsis. *Nature*, **405**, 462-466.
- 961 Park D.H., Somers D.E., Kim Y.S., Choy Y.H., Lim H.K., Soh M.S., Kim H.J., Kay S.A. & Nam H.G. (1999)
962 Control of Circadian Rhythms and Photoperiodic Flowering by the Arabidopsis GIGANTEA
963 Gene. *Science*, **285**, 1579-1582.
- 964 Pokhilko A., Hodge S.K., Stratford K., Knox K., Edwards K.D., Thomson A.W., Mizuno T. & Millar A.J.
965 (2010) Data assimilation constrains new connections and components in a complex,
966 eukaryotic circadian clock model. *Molecular Systems Biology*, **6**.
- 967 Pyl E.T., Piques M., Ivakov A., Schulze W., Ishihara H., Stitt M. & Sulpice R. (2012) Metabolism and
968 Growth in Arabidopsis Depend on the Daytime Temperature but Are Temperature-
969 Compensated against Cool Nights. *The Plant Cell*, **24**, 2443-2469.
- 970 Qin M.M., Kuhn R., Moran S. & Quail P.H. (1997) Overexpressed phytochrome C has similar
971 photosensory specificity to phytochrome B but a distinctive capacity to enhance primary leaf
972 expansion. *Plant Journal*, **12**, 1163-1172.
- 973 Quail P., Boylan M., Parks B., Short T., Xu Y. & Wagner D. (1995) Phytochromes: photosensory
974 perception and signal transduction. *Science*, **268**, 675-680.
- 975 Raschke A., Ibañez C., Ullrich K.K., Anwer M.U., Becker S., Glöckner A., Trenner J., Denk K., Saal B.,
976 Sun X., Ni M., Davis S.J., Delker C. & Quint M. (2015) Natural variants of ELF3 affect
977 thermomorphogenesis by transcriptionally modulating PIF4-dependent auxin response
978 genes. *BMC Plant Biology*, **15**, 197.
- 979 Reed J.W., Nagpal P., Bastow R.M., Solomon K.S., Dowson-Day M.J., Elumalai R.P. & Millar A.J. (2000)
980 Independent Action of ELF3 and phyB to Control Hypocotyl Elongation and Flowering Time.
981 *Plant Physiology*, **122**, 1149-1160.
- 982 Rizzini L., Favory J.J., Cloix C., Faggionato D., O'Hara A., Kaiserli E., Baumeister R., Schäfer E., Nagy F.,
983 Jenkins G.I. & Ulm R. (2011) Perception of UV-B by the Arabidopsis UVR8 Protein. *Science*,
984 **332**, 103-106.
- 985 Roden L.C., Song H.-R., Jackson S., Morris K. & Carre I.A. (2002) Floral responses to photoperiod are
986 correlated with the timing of rhythmic expression relative to dawn and dusk in Arabidopsis.
987 *Proceedings of the National Academy of Sciences*, **99**, 13313-13318.
- 988 Rudiger W., Thummler F., Cmiel E. & Schneider S. (1983) Chromophore Structure of the
989 Physiologically Active Form (Pfr) of Phytochrome. *Proceedings of the National Academy of*
990 *Sciences of the United States of America-Biological Sciences*, **80**, 6244-6248.
- 991 Sakamoto K. & Nagatani A. (1996) Nuclear localization activity of phytochrome B. *The Plant Journal*,
992 **10**, 859-868.


- 993 Salazar J.D., Saithong T., Brown P.E., Foreman J., Locke J.C.W., Halliday K.J., Carré I.A., Rand D.A. &
994 Millar A.J. (2009) Prediction of Photoperiodic Regulators from Quantitative Gene Circuit
995 Models. *Cell*, **139**, 1170-1179.
- 996 Salomé P.A., Michael T.P., Kearns E.V., Fett-Neto A.G., Sharrock R.A. & McClung C.R. (2002) The out
997 of phase 1 Mutant Defines a Role for PHYB in Circadian Phase Control in Arabidopsis. *Plant*
998 *Physiology*, **129**, 1674-1685.
- 999 Salter M.G., Franklin K.A. & Whitelam G.C. (2003) Gating of the rapid shade-avoidance response by
1000 the circadian clock in plants. *Nature*, **426**, 680-683.
- 1001 Sanchez A., Shin J. & Davis S.J. (2011) Abiotic stress and the plant circadian clock. *Plant Signaling &*
1002 *Behavior*, **6**, 223-231.
- 1003 Sanchez-Villarreal A., Shin J., Bujdoso N., Obata T., Neumann U., Du S., Ding Z., Davis A.M., Shindo T.,
1004 Schmelzer E., Sulpice R., Nunes-Nesi A., Stitt M., Fernie A.R. & Davis S.J. (2013) TIME FOR
1005 COFFEE is an essential component in the maintenance of metabolic homeostasis in
1006 Arabidopsis thaliana. *The Plant Journal*, **76**, 188-200.
- 1007 Schaffer R., Ramsay N., Samach A., Corden S., Putterill J., Carré I.A. & Coupland G. (1998) The late
1008 elongated hypocotyl Mutation of Arabidopsis Disrupts Circadian Rhythms and the
1009 Photoperiodic Control of Flowering. *Cell*, **93**, 1219-1229.
- 1010 Sellaro R., Pacín M. & Casal J.J. (2012) Diurnal Dependence of Growth Responses to Shade in
1011 Arabidopsis: Role of Hormone, Clock, and Light Signaling. *Molecular Plant*, **5**, 619-628.
- 1012 Seo H.S., Yang J.Y., Ishikawa M., Bolle C., Ballesteros M.L. & Chua N.H. (2003) LAF1 ubiquitination by
1013 COP1 controls photomorphogenesis and is stimulated by SPA1. *Nature*, **423**, 995-999.
- 1014 Sharrock R.A. & Quail P.H. (1989) Novel phytochrome sequences in Arabidopsis thaliana: structure,
1015 evolution, and differential expression of a plant regulatory photoreceptor family. *Genes &*
1016 *Development*, **3**, 1745-1757.
- 1017 Shearman L.P., Sriram S., Weaver D.R., Maywood E.S., Chaves I., Zheng B., Kume K., Lee C.C., van der
1018 G.T.J., Horst, Hastings M.H. & Reppert S.M. (2000) Interacting Molecular Loops in the
1019 Mammalian Circadian Clock. *Science*, **288**, 1013-1019.
- 1020 Sheerin D.J., Menon C., zur Oven-Krockhaus S., Enderle B., Zhu L., Johnen P., Schleifenbaum F.,
1021 Stierhof Y.D., Huq E. & Hiltbrunner A. (2015) Light-Activated Phytochrome A and B Interact
1022 with Members of the SPA Family to Promote Photomorphogenesis in Arabidopsis by
1023 Reorganizing the COP1/SPA Complex. *The Plant Cell*, **27**, 189-201.
- 1024 Shin J., Anwer M.U. & Davis S.J. (2013) Phytochrome-Interacting Factors (PIFs) as Bridges between
1025 Environmental Signals and the Circadian Clock: Diurnal Regulation of Growth and
1026 Development. *Molecular Plant*, **6**, 592-595.
- 1027 Shin J., Heidrich K., Sanchez-Villarreal A., Parker J.E. & Davis S.J. (2012) TIME FOR COFFEE Represses
1028 Accumulation of the MYC2 Transcription Factor to Provide Time-of-Day Regulation of
1029 Jasmonate Signaling in Arabidopsis. *Plant Cell*, **24**, 2470-2482.
- 1030 Shin J., Sánchez-Villarreal A., Davis A.M., Du S., Berendzen K.W., Koncz C., Ding Z., Li C. & Davis S.J.
1031 (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-
1032 dependent manner. *Plant, Cell & Environment*, **40**, 997-1008.
- 1033 Smith H. (1995) Physiological and Ecological Function within the Phytochrome Family. *Annual Review*
1034 *of Plant Physiology and Plant Molecular Biology*, **46**, 289-315.
- 1035 Somers D.E., Devlin P.F. & Kay S.A. (1998a) Phytochromes and Cryptochromes in the Entrainment of
1036 the Arabidopsis Circadian Clock. *Science*, **282**, 1488-1490.
- 1037 Somers D.E., Webb A.A., Pearson M. & Kay S.A. (1998b) The short-period mutant, toc1-1, alters
1038 circadian clock regulation of multiple outputs throughout development in Arabidopsis
1039 thaliana. *Development*, **125**, 485-494.
- 1040 Song Y.H., Smith R.W., To B.J., Millar A.J. & Imaizumi T. (2012) FKF1 Conveys Timing Information for
1041 CONSTANS Stabilization in Photoperiodic Flowering. *Science*, **336**, 1045-1049.

- 1042 Soy J., Leivar P., González-Schain N., Sentandreu M., Prat S., Quail P.H. & Monte E. (2012)
1043 Phytochrome-imposed oscillations in PIF3 protein abundance regulate hypocotyl growth
1044 under diurnal light/dark conditions in Arabidopsis. *The Plant Journal*, **71**, 390-401.
- 1045 Soy J., Leivar P., González -Schain N., Martin G., Diaz C., Sentandreu M., Al-Sady B., Quail P. H.,
1046 Monte E. (2016) Molecular convergence of clock and photosensory pathways through PIF3–
1047 TOC1 interaction and co-occupancy of target promoters. *Proceedings of the National*
1048 *Academy of Sciences of the United States of America*, **113** 4870-4875.
- 1049 Stracke R., Favory J.J., Gruber H., Bartelniewoehner L., Bartels S., Binkert M., Funk M., Weisshaar B.
1050 & Ulm R. (2010) The Arabidopsis bZIP transcription factor HY5 regulates expression of the
1051 PFG1/MYB12 gene in response to light and ultraviolet-B radiation. *Plant, Cell & Environment*,
1052 **33**, 88-103.
- 1053 Strayer C., Oyama T., Schultz T.F., Raman R., Somers D.E., Más P., Panda S., Kreps J.A. & Kay S.A.
1054 (2000) Cloning of the Arabidopsis Clock Gene TOC1, an Autoregulatory Response Regulator
1055 Homolog. *Science*, **289**, 768-771.
- 1056 Takase M., Mizoguchi T., Kozuka T. & Tsukaya H. (2013) The unique function of the Arabidopsis
1057 circadian clock gene PRR5 in the regulation of shade avoidance response. *Plant Signaling &*
1058 *Behavior*, **8**, e23534.
- 1059 Thines B. & Harmon F.G. (2010) Ambient temperature response establishes ELF3 as a required
1060 component of the core Arabidopsis circadian clock. *Proceedings of the National Academy of*
1061 *Sciences of the United States of America*, **107**, 3257-3262.
- 1062 Toth R., Kevei E., Hall A., Millar A.J., Nagy F. & Kozma-Bognar L. (2001) Circadian clock-regulated
1063 expression of phytochrome and cryptochrome genes in Arabidopsis. *Plant Physiology*, **127**,
1064 1607-1616.
- 1065 Tóth R., Kevei É., Hall A., Millar A.J., Nagy F. & Kozma-Bognár L. (2001) Circadian Clock-Regulated
1066 Expression of Phytochrome and Cryptochrome Genes in Arabidopsis. *Plant Physiology*, **127**,
1067 1607-1616.
- 1068 Troein C., Locke J.C.W., Turner M.S. & Millar A.J. (2009) Weather and Seasons Together Demand
1069 Complex Biological Clocks. *Current Biology*, **19**, 1961-1964.
- 1070 Undurraga S.F., Press M.O., Legendre M., Bujdosó N., Bale J., Wang H., Davis S.J., Verstrepen K.J. &
1071 Queitsch C. (2012) Background-dependent effects of polyglutamine variation in the
1072 Arabidopsis thaliana gene ELF3. *Proceedings of the National Academy of Sciences of the*
1073 *United States of America*, **109**, 19363-19367.
- 1074 Walters R.G., Rogers J.J.M., Shephard F. & Horton P. (1999) Acclimation of Arabidopsis thaliana to
1075 the light environment: the role of photoreceptors. *Planta*, **209**, 517-527.
- 1076 Wang H., Ma L.G., Li J.M., Zhao H.Y. & Deng X.W. (2001) Direct interaction of Arabidopsis
1077 Cryptochromes with COP1 in Light Control Development. *Science*, **294**, 154-158.
- 1078 Wang W., Barnaby J.Y., Tada Y., Li H., Tor M., Caldelari D., Lee D.U., Fu X.D. & Dong X. (2011) Timing
1079 of plant immune responses by a central circadian regulator. *Nature*, **470**, 110-114.
- 1080 Wang Z.Y. & Tobin E.M. (1998) Constitutive Expression of the CIRCADIAN CLOCK ASSOCIATED 1
1081 (CCA1) Gene Disrupts Circadian Rhythms and Suppresses Its Own Expression. *Cell*, **93**, 1207-
1082 1217.
- 1083 Wenden B., Kozma-Bognar L., Edwards K.D., Hall A.J., Locke J.C. & Millar A.J. (2011) Light inputs
1084 shape the Arabidopsis circadian system. *Plant J*, **66**, 480-491.
- 1085 Weston E., Thorogood K., Vinti G. & López-Juez E. (2000) Light quantity controls leaf-cell and
1086 chloroplast development in Arabidopsis thaliana wild type and blue-light-perception
1087 mutants. *Planta*, **211**, 807-815.
- 1088 Yang H.Q., Wu Y.J., Tang R.H., Liu D., Liu Y. & Cashmore A.R. (2000) The C Termini of Arabidopsis
1089 Cryptochromes Mediate a Constitutive Light Response. *Cell*, **103**, 815-827.
- 1090 Yang J., Lin R., Sullivan J., Hoecker U., Liu B., Xu L., Deng X.W. & Wang H. (2005) Light Regulates
1091 COP1-Mediated Degradation of HFR1, a Transcription Factor Essential for Light Signaling in
1092 Arabidopsis. *The Plant Cell*, **17**, 804-821.

- 1093 Yanovsky M.J., Mazzella M.A. & Casal J.J. (2000) A quadruple photoreceptor mutant still keeps track
1094 of time. *Current Biology*, **10**, 1013-1015.
- 1095 Yi C. & Deng X.W. (2005) COP1 – from plant photomorphogenesis to mammalian tumorigenesis.
1096 *Trends in Cell Biology*, **15**, 618-625.
- 1097 Yu J.W., Rubio V., Lee N.Y., Bai S., Lee S.Y., Kim S.S., Liu L., Zhang Y., Irigoyen M.L., Sullivan J.A., Zhang
1098 Y., Lee I., Xie Q., Paek N.C. & Deng X.W. (2008) COP1 and ELF3 Control Circadian Function
1099 and Photoperiodic Flowering by Regulating GI Stability. *Molecular Cell*, **32**, 617-630.
- 1100 Zagotta M.T., Hicks K.A., Jacobs C.I., Young J.C., Hangarter R.P. & MeeksWagner D.R. (1996) The
1101 Arabidopsis ELF3 gene regulates vegetative photomorphogenesis and the photoperiodic
1102 induction of flowering. *Plant Journal*, **10**, 691-702.
- 1103 Zeilinger M.N., Farré E.M., Taylor S.R., Kay S.A. & Doyle F.J. (2006) A novel computational model of
1104 the circadian clock in Arabidopsis that incorporates PRR7 and PRR9. *Molecular Systems
1105 Biology*, **2**.
- 1106 Zhu D., Maier A., Lee J.H., Laubinger S., Saijo Y., Wang H., Qu L.J., Hoecker U. & Deng X.W. (2008)
1107 Biochemical Characterization of Arabidopsis Complexes Containing CONSTITUTIVELY
1108 PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA Proteins in Light Control of Plant
1109 Development. *The Plant Cell*, **20**, 2307-2323.
- 1110 Zuo Z., Liu H., Liu B., Liu X. & Lin C. (2011) Blue Light-Dependent Interaction of CRY2 with SPA1
1111 Regulates COP1 activity and Floral Initiation in Arabidopsis. *Current biology : CB*, **21**, 841-847.







Shining a light on the Arabidopsis circadian clock

Rachael J. Oakenfull, Seth J. Davis

Department of Biology, University of York, York, UK

Corresponding author: Seth J. Davis

E-mail: seth.davis@york.ac.uk

For Review Only

14 Abstract

15
16 The circadian clock provides essential timing information to ensure optimal growth to prevailing
17 external environmental conditions. A major time-setting mechanism (*zeitgeber*) in clock
18 synchronisation is light. Differing light wavelengths, intensities and photoperiodic duration are
19 processed for the clock-setting mechanism. Many studies on ~~the~~ light-input pathways to the clock
20 have focused on *Arabidopsis thaliana*. Photoreceptors are specific chromic proteins that detect light
21 signals and transmit this information to the central circadian oscillator through a number of different
22 signalling mechanisms. The most ~~well characterised~~ well characterised clock-mediating
23 photoreceptors are cryptochromes and phytochromes, detecting blue, red and far-red wavelengths
24 of light. Ultraviolet and shaded light ~~are also also are~~ processed signals to the oscillator. Notably, the
25 clock reciprocally generates rhythms of photoreceptor action leading to so-called gating of light
26 responses. ~~A number of~~ intermediate proteins, such as Phytochrome interacting factors (PIFs),
27 constitutive photomorphogenic 1 (COP1) and EARLY FLOWERING 3 (ELF3), have been established in
28 signalling pathways downstream of photoreceptor activation. However, the precise details for these
29 signalling mechanisms are not fully established. This review highlights both historical and recent
30 efforts made to understand overall light input to the oscillator, first looking at how each wavelength
31 of light is detected, this is then related to known input mechanisms and their interactions.
32

Review Only

The circadian clock

The circadian clock allows plants as sessile organisms to synchronise with diurnal changes in the environment (Dodd *et al.*, 2005). Daily external environmental stimuli are required to initiate circadian oscillations and to maintain synchronicity with the external environment. This process is called entrainment. The environmental cues governing these processes are termed *zeitgebers* (from German: "time givers"). The ability to synchronise with the external environment efficiently confers enhanced fitness (Michael *et al.*, 2003).

Diurnal changes in cellular processes controlled by the clock allow plants to anticipate, and therefore better survive, a range of stresses (Sanchez *et al.*, 2011). Diurnal changes have been shown to occur in cold/freezing tolerance (Fornara *et al.*, 2015, Nakamichi *et al.*, 2009), drought tolerance (Habte *et al.*, 2014), pathogen response (Shin *et al.*, 2012, Wang *et al.*, 2011) and photosynthesis (Pyl *et al.*, 2012). This synchronisation is the product of a large number of rhythmically regulated cellular processes (Bujdosó & Davis, 2013, Hanano *et al.*, 2008), many of which are triggered by light perception (Wenden *et al.*, 2011). Perception of daily *zeitgebers*, such as changes in light and temperature (Chow *et al.*, 2014, Harmer, 2009), enable plants to reset the clock at dawn, and feed back to the central oscillator. For this light entrainment, photoreceptors play a major role (Somers *et al.*, 1998a, Toth *et al.*, 2001). The circadian clock was derived from the principle of inter-connected, positive and negative feedback loops (Shearman *et al.*, 2000). For the purpose of this review, light input into the *Arabidopsis thaliana* (*Arabidopsis*) circadian clock will be the focus of discussion, with a brief examination of clock components.

In *Arabidopsis*, morning expressed Myb-like transcription factors *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) (Wang & Tobin, 1998) and *LATE ELONGATED HYPOCOTYL* (*LHY*) (Schaffer *et al.*, 1998) antagonize expression of the evening expressed pseudo-response regulator (PRR) *TIMING OF CAB EXPRESSION 1* (*TOC1*) (Strayer *et al.*, 2000). These three genes form the core negative feedback loop of the circadian oscillator (Alabadi *et al.*, 2001, Gendron *et al.*, 2012) (Figure 1). Several other genes form additional loops within this core oscillator. In day time *CCA1* and *LHY* repress expression of the pseudo-response regulators *PRR5*, *PRR7* and *PRR9* (Adams *et al.*, 2015, Kamioka *et al.*, 2016), as well as *TOC1*, *GI*, and the genes that generate the evening complex (Locke *et al.*, 2006, Nakamichi *et al.*, 2009, Pokhilko *et al.*, 2010, Zeilinger *et al.*, 2006). *GIGANTEA* (*GI*) is evening expressed and is proposed to form an additional negative feedback-loop with *TOC1* (Locke *et al.*, 2006). All of these loops are connected through the action of the evening complex formed by *LUX ARRHYTHMO* (*LUX*), *EARLY FLOWERING 3* (*ELF3*) and *EARLY FLOWERING 4* (*ELF4*), which directly inhibits the expression of *PRR9* (Helfer *et al.*, 2011, Herrero *et al.*, 2012), *PRR7*, *GI* and *LUX* (Mizuno *et al.*, 2014). The absence of even one component of the evening complex gives rise to plants that are photoperiod insensitive. This results in early flowering, long hypocotyl growth and arrhythmicity of the free-running circadian period (Hazen *et al.*, 2005, McWatters *et al.*, 2007, Onai & Ishiura, 2005, Thines & Harmon, 2010). The importance of the three evening-complex components is thusly highlighted in maintaining a functional circadian clock, and therefore the physiological processes controlled by the clock, such as the input of diurnal photoperiod information, (Covington *et al.*, 2001, Más *et al.*, 2003, Mizoguchi *et al.*, 2005, Park *et al.*, 1999).

Effects of light on the clock

Light changes throughout a day-night cycle are pronounced and thus robustly entrain the clock. In the light phase of a daily cycle, the dark to light transition of dawn is used as a time setting checkpoint (Millar *et al.*, 1995). Prolonged darkness causes many of the core genes in the *Arabidopsis* central oscillator to rapidly become arrhythmic, due to the lack of essential light time setting cues (Figure 2a) (Millar *et al.*, 1995). This dampening effect, leading to arrhythmicity is particularly noticeable in the absence of media containing sucrose. In prolonged darkness, sucrose can act as a substitute for light in maintaining rhythmicity for a number of days (Bläsing *et al.*, 2005). Light has two main modes to set the clock. The first is parametric entrainment; gradual entrainment of the clock, such as the acceleration of the clock induced by increased light perception, which

eventually leads to a phase shift of the clock back to a correct resonance. Parametric entrainment follows Aschoff's rule, as light intensity increases, the speed of the clock increases. As intensity decreases, the speed of the clock slows (Aschoff, 1979), (Figure 2B/C). Increases in light intensity, lead to decreases in periodicity (Somers *et al.*, 1998a). The second light-induced time-setting mechanism is non-parametric entrainment: rapid re-entrainment. This leads to a rapid time setting of the clock at dawn (Millar & Kay, 1996). Non-parametric entrainment requires an extended light exposure far beyond that which activates light-regulated gene expression (Millar & Kay, 1996). Metabolic entrainment is also a mechanism for non-parametric entrainment (Haydon *et al.*, 2013, Sanchez-Villarreal *et al.*, 2013; Haydon & Webb, 2016; Shin *et al.*, 2017). The different photoreceptors and photochromic proteins involved in light entrainment are described in more detail below.

How are different wavelengths of light input to the *Arabidopsis* clock?

Diurnal organisms, particularly plants, are subjected to Aschoff's rule: an increase in light intensity accelerates the circadian-oscillator speed leading to shortening of periodicity (Aschoff, 1979). Light input to the circadian clock is presumed to occur through the action of different types of photoreceptors (Somers *et al.*, 1998a). There are more than ten known circadian-associated photoreceptors (Edwards *et al.*, 2015). These can be split into four classes: phytochromes, cryptochromes, ZTL/FKF1/LKP2 family, and UVR8. Each receptor contributes in the dose-dependent perception of far-red, red, blue, and ultra-violet light (Cashmore *et al.*, 1999, Mas *et al.*, 2000, Rizzini *et al.*, 2011, Song *et al.*, 2012). It is presumed that the input of this information is co-ordinately relayed to the central oscillator.

Both phytochromes and cryptochromes play key roles in light responsive time setting mechanisms, in a manner that follows Aschoff's rule (Devlin & Kay, 2000a, Somers *et al.*, 1998a). This is due to the ability of both phytochromes and cryptochromes to form photoreceptor complexes (Más *et al.*, 2003) that are genetically interactive in clock function (Devlin & Kay, 2000a). Excitation of these photoreceptors cause the central oscillator to accelerate, changing the overall speed of the clock (Devlin & Kay, 2000b, Herrero *et al.*, 2012, Kolmos *et al.*, 2011, Somers *et al.*, 1998a, Somers *et al.*, 1998b). There are a number of different known mechanisms through which light absorption by photoreceptors input environmental information to the oscillator, however, these mechanistic details are not complete. Regulation of transcription by circadian gating restricts changes in RNA levels to specific times of day. Therefore preventing transcription of some light-regulated clock genes in response to unexpected external stimuli, for example light pulses during the night (Millar & Kay, 1996). Light regulation of myb transcription factors, such as CCA1 and LHY effect the transcription and stability of other clock components, such as PRR9/7 (Carre & Kay, 1995). Messengers such as Ca²⁺ and calmodulin signalling may also affect circadian regulation in response to light (Johnson *et al.*, 1995, Millar & Kay, 1996). Light also directly controls the degradation of PRR5, PRR7, PRR9, TOC1 and GI proteins (David *et al.*, 2006, Farré and Kay, 2007, Ito *et al.*, 2007, Kiba *et al.*, 2007, Más *et al.*, 2003, Matsushika *et al.*, 2000). These degradation events then act on outputs within a diurnal context, which change in duration throughout the season (Davis, 2002, Guerriero *et al.*, 2012, Salazar *et al.*, 2009, Song *et al.*, 2012, Troein *et al.*, 2009). Light thus has multiple mechanistic inputs to clock processes, all of which control entrainment. How each individual wavelength of light is input to the clock will be discussed below.

Red light

Phytochromes are predominantly red-light photoreceptors, absorbing maximally at wavelengths between 600 and 700nm (Somers *et al.*, 1998a). *Arabidopsis* has five phytochromes (Sharrock & Quail, 1989), phyA-phyE (Mathews & Sharrock, 1997). Each phytochrome acts as a light input sensor to form regulatory feedback loops within the circadian clock. Phytochromes are in turn reported to be negatively regulated by the clock through cryptochrome (CRY) signals (Devlin & Kay, 2000a, Mas *et al.*, 2000). Phytochromes exist in two interconvertible forms; the inactive Pr form is

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

converted by red light to the active Pfr form which can be converted back to the inactive Pr state by far-red light (Rudiger *et al.*, 1983). These conversion events between active and inactive forms of phytochrome are essential to light input to the clock, as discussed below (see far red, PIFs, ELF3). Each of the five phytochromes play distinct roles in light sensing.

phyA mediates entrainment responses to low intensity red light and pulses of light (Quail *et al.*, 1995, Somers *et al.*, 1998a). A *PHYA* deficiency mutation, results in an altered period length in dim red light (Somers *et al.*, 1998a). It is not known how phyA signals to the clock as it has not been reported to directly bind to a clock-associated factor, in contrast to the other four phytochromes (Huang *et al.*, 2016).

phyB is the main detector for high intensity red light (Somers *et al.*, 1998a). Both phyB and phyD are able to detect red and far-red wavelengths of light (Aukerman *et al.*, 1997, Devlin *et al.*, 1999). phyB physically interacts with ELF3 in the central oscillator to provide a direct light input to the clock (Kolmos *et al.*, 2011, Liu *et al.*, 2001). *phyb* mutants show an altered response to shade avoidance (Smith, 1995), which is also a phenotype of the *elf3* mutant (Huang *et al.*, 2016). phyC to phyE also interact with ELF3 protein (Huang *et al.*, 2016), but this has not yet been connected to the clock (Liu *et al.*, 2001). Under high fluence red light, *phyb* mutants and the phyB overexpressor have a period defects and altered phase (Anderson *et al.*, 1997, Kolmos *et al.*, 2011, Salomé *et al.*, 2002, Somers *et al.*, 1998a). Also altered cryptochrome signalling (see blue light below), phyB and CRY2 physically interact by translocating to the nucleus in red light (Mas *et al.*, 2000), where phyB is then suppressed by CRY2 (Mas *et al.*, 2000). This alters clock performance under white light conditions (red and blue light together) (Devlin & Kay, 2000a).

In non-peer-reviewed work, phyC was found to play a role in white-light input and red-light detection. Mutations in *PHYC* result in a long-period phenotype, which was shown to be temperature dependant, suggesting that phyC inputs not only light information to the clock, but also temperature (Edwards *et al.*, 2015, Franklin *et al.*, 2003, Qin *et al.*, 1997). phyE along with phyD plays a role in controlling the period length of *CAB* gene expression, however, many of the clock effects of phyE and phyD are masked by phyB (Franklin & Quail, 2010). phyE works with phyB and phyD in the regulation of shade avoidance (Devlin *et al.*, 1998). Interestingly, the promoters of *PHYA* and *PHYB* are down-regulated by light, whereas the *PHYC* promoter is upregulated (Tóth *et al.*, 2001), *PHYD* and *PHYE* do not show changes in expression in response to light changes. *PHYB,D,E* mediate high fluence red light input to the clock with *PHYA,B,D,E* acting additively to input red-light information to the clock, as a result the clock runs faster as the detected intensity of red light increases (Devlin & Kay, 2000a). The absence of all five phytochromes results in severally attenuated rhythms, but not a total loss of clock function (Hu *et al.*, 2013). Together, all five phytochromes play differing roles in mediating light-dependant changes in periodicity.

Blue light

Cryptochromes are blue light (492 to 455nm) and UVA photoreceptors present in both plants and animals (Cashmore *et al.*, 1999). The *HY4* locus was found to encode cryptochrome 1 (CRY1). It was identified due to *cry1* (*hy4*) mutants growing with a long-hypocotyl phenotype and being unable to respond to blue light (Ahmad & Cashmore, 1993, Koornneef *et al.*, 1980), *cry1/hy4* plants are also late flowering (Goto *et al.*, 1991, Millar *et al.*, 1995). *cry1* mutants have a long period under blue light (Somers *et al.*, 1998a), suggesting CRY1 acts as a photoreceptor for blue-light entrainment of the clock (Devlin & Kay, 2000a). Overexpression of *CRY1* caused increased sensitivity to blue light and period shortening (Lin *et al.*, 1996, Somers *et al.*, 1998a). CRY1 is a soluble protein when grown in both light and dark conditions in *Arabidopsis* (Lin *et al.*, 1996), CRY1 is more stable than CRY2 and works at higher light intensities (Lin *et al.*, 1998). Cryptochrome 2 (CRY2) can detect low intensity light and is rapidly degraded under blue light (Lin *et al.*, 1998). In light, *CRY2* promoter activity is down-regulated whereas, *CRY1* is upregulated (Tóth *et al.*, 2001). The *cry2* mutation alters sensitivity to photoperiod and flowering in *Arabidopsis*, but does not have a detectable individual effect on circadian rhythm (Devlin & Kay, 2000a, Guo *et al.*, 1998). Overexpression of either *CRY1* or

CRY2 gives rise to a higher blue-light sensitivity under low light conditions than in the individual overexpression lines (Ahmad *et al.*, 1998a). Double mutant *cry1, cry2* plants are rhythmic, suggesting that although CRY1 inputs blue light into the clock CRY1 and 2 are not part of the central oscillator (Devlin & Kay, 2000a). However, CRY1 and CRY2 gene expression oscillates with a circadian rhythm under constant light (Harmer *et al.*, 2000). CRY1 and 2 work together to input information to the clock in a similar way to phyA and B, but at differing light intensities.

Phytochromes are able to absorb low fluence blue light alongside CRY1 for period length control. *phyA* mutants show a period lengthening effect when free run under blue light (Somers *et al.*, 1998a). Without phyA detection of blue light, the input relies on CRY1 alone causing the period to lengthen as the plant detects less light than the actual ambient intensity of irradiation. Conversely, *PHYA* overexpression has been proposed to cause period shortening under blue light, as more blue light is processed as an input than the actual ambient light intensity. Phytochromes thus also work in blue-light signalling to the clock.

High light synergism - (White light)

White light comprises of multiple light wavelengths. As such, interactions between phytochromes and cryptochromes are needed to input this information into the circadian clock. These interactions were found with loss of function mutants for both phytochromes and cryptochromes. In wild-type plants *CAB2* period decreases as light intensity increases, *cry2* mutants were found to be deficient in a white light response as they have a *CAB2* period increase in response to high light (Mas *et al.*, 2000). This period increase was not detected in either red or blue light alone, suggesting that to be active CRY2 needs multiple wavelengths of light simultaneously and phyB (Mas *et al.*, 2000). CRY1 was also found to be required for phyA signalling as *cryCRY1* and *cryCRY2* mutants are unable to detect red light above the fluence range of both phyA and phyB (Devlin & Kay, 2000a). Light induces nuclear compartmentalisation of phytochromes where phyA and phyB directly interact with CRY1 and CRY2 (Mas *et al.*, 2000), the kinase activity of phyA phosphorylates CRY1 and CRY2 (Ahmad *et al.*, 1998b).

Phytochromes and cryptochromes facilitate signal integration of multiple light cues. CRY2 is activated when illuminated by multiple wavelengths of light suggesting it is needed for phytochrome activation (Mas *et al.*, 2000). This is also highlighted as both cryptochromes reach peak RNA expression with a similar expression pattern to the corresponding phytochromes (Toth *et al.*, 2001). It was found that the active Pfr form of phytochrome is needed for CRY2 expression, CRY2 then suppresses *PHYB* expression. However, PfrB is able to override CRY2 signalling to flowering time control via pathways, such as COP1 (see below). PfrB binding to the intermediate SPA1 allows degradation of the COP1-SPA1 complex, which is needed as an intermediate of CRY1/2 induced inhibition of photomorphogenic factors such as HFR and CO (Mas *et al.*, 2000, Sheerin *et al.*, 2015). Additionally, both CRY1 and CRY2 were found to be phosphorylated by the kinase activity of phyA (Ahmad *et al.*, 1998b; Mas *et al.*, 2000). It could be considered that phytochromes and cryptochromes work together in the "white light" response, which is a more than the additive effect of plants grown under blue and red light. However, quadruple mutants for *phyA, phyB, cry1* and *cry2* still showed rhythmic leaf movement in response to light-dark cycles suggesting that other photoreceptors must play a role in overall light input to the circadian clock (Yanovsky *et al.*, 2000). However, the exact relationship between phytochromes and cryptochromes is yet to be resolved.

ZTL family; blue-light absorbing with action under red-light and darkness

Zeitlupe (ZTL), also reported as ADAGIO1 (ADO1), links light input by both cryptochromes and phytochromes to the clock (Jarillo *et al.*, 2001, Kim *et al.*, 2007). *ZTL* mutant lines showed altered cotyledon movement and gene expression under different light conditions (Jarillo *et al.*, 2001). Under blue and white light, *ztl* plants have a long period whereas under red light the *ztl* lines were reported to be arrhythmic for *CCR2* expression, cotyledon movement and stem elongation (Jarillo *et al.*, 2001). *ZTL* mutants were found to have a long-period phenotype for *CAB/TOC1* under

Formatted: Font: Italic

red light (Kevei *et al.*, 2006, Kim *et al.*, 2005). ZTL thus is required for the perception of multiple wavelengths of light into the oscillator.

ZTL encodes a protein reported to be a blue-light photoreceptor, as it contains a PAS domain, F box domain linking proteins to a SCF ubiquitination complex, kelch repeats and a light, oxygen or voltage (LOV) domain allowing protein-protein interactions (Mas *et al.*, 2003). Interactions between TOC1 and ZTL were found to occur through these kelch-repeat zones (Kevei *et al.*, 2006). The PAS/ LOV domain were identified as essential for coupling ZTL to red light (Kevei *et al.*, 2006), which was then found to occur through ZTL binding to the C-terminus of PhyB and CRY1 (Kim *et al.*, 2007).

ZTL is constitutively expressed at the RNA level, however, oscillations in ZTL protein levels are seen (Kim *et al.*, 2007). These are proposed to result from the binding of Gigantea (GI) to maintain the stability of ZTL. ZTL protein folding is chaperoned by HSP90 (Kim *et al.*, 2011), GI binds to the ZTL-HSP90 complex to ensure specificity of protein folding (Cha *et al.*, 2017, Kim *et al.*, 2011). Interactions between ZTL and GI are enhanced by blue light through the LOV domain in ZTL (Kim *et al.*, 2007). ZTL controls proteasomal degradation of TOC1 (Más *et al.*, 2003). This ZTL-GI interaction is believed to control a central part of the circadian oscillator. ZTL and ELF3 were reported to have opposite effects on clock function. *ztl* mutants and *ELF3* overexpression lines show a lengthened circadian period in light. Conversely *elf3* mutants and ZTL overexpression lines are reported as arrhythmic under constant light (LL) (Kim *et al.*, 2005). However, the *elf3-ztl* double mutant showed that ELF3 and ZTL have additive effects on the clock (Kim *et al.*, 2005). As GI controls the HSP90 mediated stabilisation of ZTL protein (Cha *et al.*, 2017, Kim *et al.*, 2011), ZTL protein then causes protein depletion of TOC1 via ubiquitination (Kim *et al.*, 2011). ELF3 interacts as a substrate adaptor for COP1 (an E3 ubiquitin ligase) to bind to and degrade GI protein, as a light input signal and indicator of day length in response to CRY2 (Yu *et al.*, 2008). The reduction of GI then prevents the formation of stable ZTL protein. ~~Therefore Consequently preventing this prevents~~ ZTL-mediated inhibition of TOC1 in the central oscillator, and ~~facilitating facilitates~~ TOC1 action. ~~ZTL also negatively regulates PRR5 by targeting PRR5 protein for degradation by via the 26S proteasome (Fujiwara et al., 2008, Kiba et al., 2007). As PRR5 forms a negative regulatory feedback loop with LHY/ CCA1, ZTL also therefore indirectly plays a role in the regulation of LHY/CCA1 within the central oscillator (Baudry et al., 2010). The relationship of ZTL to other clock components is yet to be established.~~

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Far-red light

phyA is the presumed photoreceptor for detecting monochromatic far-red light. Mutations in *PHYA* resulted in loss of capacity for clock function (Wenden *et al.*, 2011). ELF4 was proposed to restrict far-red perception in those studies. Interestingly, the active form of phyA (phyA-Pfr) is formed under far-red light (Clough & Vierstra, 1997), given that far red converts the Pfr form of phytochrome back to the inactive Pr form. In part perhaps phyA evolved the ability to form Pfr under far red as a response to the change in light quality at the end of the day, which signals the transition from day to night and therefore the associated changes in environment. However, far red can also be a signal of shade due to far red being one of the only wavelengths of light able to pass through leaves (Federer & Tanner, 1966), suggesting that there may be different mechanisms to entrain the clock in these two different circumstances, as described in the next section. Plants in constant far-red light have a faster clock and show high expression of evening genes, such as *PRR1/TOC1*, and low expression of the morning genes *CCA1* and *LHY* (Wenden *et al.*, 2011). The exact mechanism of far-red input to the clock is not fully characterised. However, far red has been shown to be involved in the prevention of the interaction between Pfr and PIF3 (Martínez-García *et al.*, 2000). ELF4 was identified as playing a role in mediating far-red light input to the clock (Wenden *et al.*, 2011), Far-red light was used to aid recovery of rhythmicity in the otherwise arrhythmic *elf3* and *elf4* mutants (Kolmos *et al.*, 2011, Wenden *et al.*, 2011).

Shaded light

White light with supplementary far-red light causes the clock to slow down (Jiménez-Gómez *et al.*, 2010). Under shade, far red and potentially green light are present; there is a large overlap between far-red signalling and shade. Shade however, is a useful environmental indicator to plants for neighbour detection. *phyA* is thought to have the most involvement in mediating far-red signalling, but *phyB* also plays a key role (Kolmos *et al.*, 2011, Wenden *et al.*, 2011). Shading plants during the afternoon was found to have the greatest effect (Sellaro *et al.*, 2012). Responses to shade involve the degradation of phytochrome interacting factors, namely PIF4 and 5 (Lorrain *et al.*, 2008). PRR5 was found to regulate the shade-avoidance response by controlling PIF4 and PIF5, as well as downstream components of the phytochrome-mediated signalling pathway. Furthermore ZTL induces degradation of PRR5. However, this degradation was found to be repressed under blue light. It was suggested that PRR5 gates phytochrome mediated shade responses (Takase *et al.*, 2013). *ELF3* and *LUX* mutants (both components of the evening complex) show a reduced response to all wavelengths of light therefore growing with elongated hypocotyls as though under shade (Jiménez-Gómez *et al.*, 2010, Sellaro *et al.*, 2012, Zagotta *et al.*, 1996). This implies that *ELF3* and the evening-complex also play a role in the shade response to the clock (Kolmos *et al.*, 2011).

UV-B light

Ultraviolet B light (UVB) can be one of the more damaging wavelengths present in sunlight. UVB is a wavelength that is easily absorbed and damages both DNA and proteins (Jansen *et al.*, 1998), thus making UVB a useful light signal, but at the cost of inducing a stress response. UVB is an "anti-shade" signal informing a plant it is under direct sunlight. At lower-fluence rates, UV-B light is able to control development, promote photomorphogenesis, and drive gene expression (Heijde & Ulm, 2012). *Ultraviolet resistance locus 8 (UVR8)* drives signalling for the majority of UVB responses (Favory *et al.*, 2009, Rizzini *et al.*, 2011). Under UVB light, COP1 promotes the induction of elongated hypocotyl 5 (HY5) and HY5 homologue (HYH) which induce stress responses such as flavonoid biosynthesis to reduce UVB induced damage (Stracke *et al.*, 2010). UVR8 and COP1 are also crucial for UVB light entrainment of the clock (Fehér *et al.*, 2011). Under white light supplemented with UV-B light, COP1 induces HY5 and HYH, HY5 and HYH have not yet been implicated for clock entrainment by UVB (Fehér *et al.*, 2011). UVR8 is able to mediate both parametric and non-parametric entrainment, by inducing *PRR9* and *GI* under continuous light, alongside an increase in *CCA1* and *ELF3* response to UVB light pulses. UVR8 was identified as the UVB receptor that can mediate signal input to the oscillator, due to the fact that *uvr8* plants cannot input UVB light into the oscillator, (Fehér *et al.*, 2011, Heijde & Ulm, 2012). It has been proposed that UVR8 mediates UVB light input into the central oscillator by inhibiting PIF4 in the presence of UVB light. This requires COP1-mediated repression of *PIF4* transcript, and also through the stabilisation of HFR, which inhibits PIF4 (Hayes *et al.*, 2017). Canonical pathways used in UVB signalling mediate entrainment in the clock, but the critical nodes in entrainment are not fully resolved.-(Hayes *et al.*, 2017).

Green light

Many studies have been carried out to test the physiological effects that occur as a consequence of increased or absent green-light wavelengths. The mechanisms of sensing and input to the circadian clock are yet to be understood. It is thought that green wavelengths can operate via both a cryptochrome dependant and independent pathway (Folta & Maruhnich, 2007). Green light can reverse the effect of blue light on hypocotyl elongation (Bouly *et al.*, 2007, Folta, 2004), potentially due to the reversal of the blue light degradation of CRY1 (Bouly *et al.*, 2007). This could then have an effect on photoperiod and subsequently flowering time (Banerjee *et al.*, 2007, Folta & Maruhnich, 2007). The association of green light to cryptochromes was also shown by (Lin *et al.*, 1996) as overexpression of CRY1 causes increased sensitivity to green light. A reversible interaction between CRY and green light similar to that found for phytochrome in red and far red light suggests

that there are intermediate signalling factors similar to PIFs that are yet to be identified. It is thus plausible that green light could entrain the clock, but no definitive experiments have tested this.

Transcriptional regulation of photoreceptors by the clock.

The circadian clock generates rhythms of RNA and/or accumulation for all photoreceptor classes. Starting with the discovery that *phyB* mRNA is rhythmic (Bognár *et al.*, 1999, Toth *et al.*, 2001, Tóth *et al.*, 2001), subsequent findings revealed that all five phytochromes in Arabidopsis cycle. Interestingly, sub-nuclear accumulation of phytochrome holoprotein also appears to be under clock control. However, the implications of this are currently unclear. Similarly, *CRY* genes are rhythmic (Toth *et al.*, 2001). For UVR8, as UV light induces dimer disassembly to a monomer state, a diel cycle of dimers at night and monomers during the day occur (Findlay & Jenkins, 2016). *UVR8* mRNA displays robust circadian rhythms with a peak around subjective dusk (Mockler *et al.*, 2007). For ZTL, the mRNA generated does not cycle, but robust protein cycling is readily detectable. Together it is clear light receptors that act as input components to the clock, are themselves circadian-output regulated.

Phytochrome input to the central oscillator

Light input to the central oscillator is a daily *zeitgeber*, but the central oscillator also acts as a feedback mechanism to phytochromes over the day. The oscillator receives a number of light queues during the light phase of each day from photoreceptors detecting the different ratio of light wavelengths across the day. In turn, the oscillator inhibits expression of phytochrome proteins at points hypothesised in Figure 3.

At dawn (ZT0), *PHYC* peaks with the return of light and changing temperature (Toth *et al.*, 2001). HFR increases due to increased stability (Yang *et al.*, 2005). PIF4/5 expression peaks at dawn (Nomoto *et al.*, 2012), due to the lack of phytochromes, thus preventing phytochrome-induced degradation at dawn (Shin *et al.*, 2013). PIF4/5 continue to be present throughout the light phase, but are slowly degraded by interactions with the Pr form of phytochrome, PIFs can also interact with LHY, CCA1. PfrB interacts with PIF3, where PIF3 then binds to the G-box domain of *CCA1/LHY* promoters. CCA1 and LHY, as MYB transcription factors, then control other genes within the central oscillator, such as *PRR5*, 7 and 9, as well as non-circadian genes (Martínez-García *et al.*, 2000, Wang & Tobin, 1998).

PHYD and *E* are expressed two hours after dawn (Toth *et al.*, 2001). During the first half of the light phase (ZT0-6) there is an increase in light intensity up to ZT6, and alongside this, an increasing expression of phytochromes. *PHYB* and *CRY1* reach peaks in expression around mid-day (ZT6) as both *phyB* and *CRY1* work at high-light intensities (Lin *et al.*, 1996, Lin *et al.*, 1998, Toth *et al.*, 2001). *CAB1* expression peaks around mid-day as does the expression of HFR (Yang *et al.*, 2005). HFR is thought to interact with PIF3 (Fairchild *et al.*, 2000), but the mechanism through which this happens is not fully understood.

Both *PHYA* and *CRY2* peak towards the latter half of the light phase (ZT6-12) with the decreasing light and increasing far red intensity (Toth *et al.*, 2001). *CRY2* detects lower intensity light (Lin *et al.*, 1998), and the Pfr form of phytochrome is needed for *CRY2* activation (Mas *et al.*, 2000). *CRY2* then suppresses *PHYB* expression (Mas *et al.*, 2000). *phyA* is essential in controlling the clock in low light (Quail *et al.*, 1995, Somers *et al.*, 1998a) and far red conditions (Wenden *et al.*, 2011), potentially through the Pfr form of phytochrome being unable to interact with PIF3 (Martínez-García *et al.*, 2000).

At dusk, *CAB2* expression decreases, and *COP1* expression increases (Yang *et al.*, 2005), allowing *COP1* suppression of HFR throughout the dark phase of the day. *COP1* accumulation along with *ELF3* inhibits *GI* late in the afternoon (Yu *et al.*, 2008). PILs are rapidly produced during the first hour of shade, early into the dark phase and work with *TOC1* to restrict growth (Salter *et al.*, 2003).

PIF3 is at its highest level at dusk due to its interactions with the Pfr form of phytochrome and the highest level of Pfr being present just before dusk. Allowing information on high levels of far red light to be input to the central oscillator (Martínez-García *et al.*, 2000). The evening complex (ELF3, ELF4, LUX) inhibits the transcription of *PIF4/5* at dusk (Herrero *et al.*, 2012, Nusinow *et al.*, 2011; Raschke *et al.* 2015). This allows PIF protein to accumulate stably due to the lack of phytochrome inhibition overnight, which thus promotes growth, reaching a maximal level at dawn (Shin *et al.*, 2013, Delker *et al.* 2014; Raschke *et al.* 2015).

COP1

Constitutive photomorphogenic 1 (COP1) is an E3 ubiquitin ligase, mediating day length input to the clock and flowering time. COP1 is negatively regulated by a direct protein-protein interaction with CRYs (Jang *et al.*, 2008, Wang *et al.*, 2001, Yang *et al.*, 2000). phyA and B affect the nuclear abundance of COP1 (Osterlund *et al.*, 1999), as the C-terminal domain of phyB directly interacts with COP1 (Millar *et al.*, 1994). COP1 acts as an intermediate, inputting photoperiodic information from PHY and CRY into the oscillator. COP1 in turn plays a negative regulatory role targeting phyA, phyB and HFR1 for ubiquitination (Osterlund *et al.*, 2000, Seo *et al.*, 2003, Yang *et al.*, 2005).

CRY1, CRY2, phyA and phyB all interact with COP1 via Suppressor of Phytochrome A (SPA). SPA1 is a nuclear localised repressor of phytochrome signalling (Hoecker *et al.*, 1999), which interacts with COP1 (Hoecker & Quail, 2001). SPA1 contains a coiled-coil domain that enhances the E3 ligase activity of COP1 on its targets (Seo *et al.*, 2003). The interactions between the 4 known SPA proteins and COP1, negatively regulate light signalling in response to certain wavelengths of light (Laubinger *et al.*, 2004, Zhu *et al.*, 2008). COP1-SPA1 interaction is repressed by activated CRY1 in blue light (Lian *et al.*, 2011), CRY2 interacts with COP1 via SPA1 to allow COP1 proteolysis of CO to control flowering time under blue light (Zuo *et al.*, 2011). The CRY1-SPA1 interaction enhances CRY2-SPA1 activity in response to blue light to suppress COP1 activity resulting in a suppression of CO degradation (Ordoñez-Herrera *et al.*, 2015, Zuo *et al.*, 2011).

In seedlings, phyA binds to SPA1 and 2 whereas, in adult plants phyA binds to SPA3 and 4 (Laubinger *et al.*, 2004). Binding of phyB to SPA1 is Pfr dependant allowing degradation of COP1/SPA1 in light conditions that promote nuclear accumulation of phyA and B, this enhances light responses, as the disruption of COP1/SPA1 interaction prevents degradation of photomorphogenic factors such as HFR, and HY5 (Sheerin *et al.*, 2015). The COP1/SPA complex is an important factor in repression of light responses in darkness, as the COP1/SPA complex interacts directly with photoreceptors leading to its inactivation (Huang *et al.*, 2014).

Within the central oscillator, COP1 interacts with ELF3 to mediate COP1 degradation of GI late in the afternoon (Yu *et al.*, 2008), potentially using ELF3 as an adaptor for COP1 binding to GI (Liu *et al.*, 2008). It is also possible that COP1 regulates the level of ELF3 present, in *cop1* mutants ELF3 protein accumulates to higher levels than in the wild type, but the mRNA levels remain unchanged (Liu *et al.*, 2001). [The result](#)

COP1 is also involved in UVB signalling as *cop1* mutants are deficient in a UVB response (Oravec *et al.*, 2006). In the early stage of UVB signalling, UVR8 and COP1 directly interact in the nucleus (Favory *et al.*, 2009), UVR8 and COP1 were found to be essential for UVB entrainment (Fehér *et al.*, 2011). HY5 and HYH which are also important components of UVB signalling are regulated by COP1 (Brown & Jenkins, 2008). In the light COP1 detaches from HY5 allowing stabilisation and the light responsive target genes of HY5 to be activated (Yi & Deng, 2005). COP1 plays an important mediator role in the input of light from photoreceptors to the oscillator. In turn its regulation is dependent on photoreceptors, the short-period phenotype in mutant lines shows that COP1 plays a negative regulatory role on the clock.

PIFs and PILs

Phytochrome interacting factors (PIF) are a family of basic helix loop helix transcription factors. There are 4 well characterised PIFs, PIF1,3,4,5 (Leivar *et al.*, 2012). (Pfeiffer *et al.*, 2012) PIFs are unstable in the light due to their interaction with active phytochrome causing phosphorylation and subsequent degradation (Leivar *et al.*, 2012, Soy *et al.*, 2012). The most well characterised PIF is PIF3, which was found to interact with the Pfr form of phytochrome B (PfrB) acting as a bridge between PfrB and its target gene by translocating PfrB to the nucleus. Thus allowing light induced control of gene expression, as PIF3 does not interact with the Pr form of phytochrome (Martínez-García *et al.*, 2000, Pfeiffer *et al.*, 2012). PIFs are also able to input information to the clock via direct interaction with clock genes that contain a G-box motif in their promoter, PIFs can interact directly with *LHY*, *CCA1*, *PRR5*, *PRR7*, *PRR9* and *LUX* (Martínez-García *et al.*, 2000). This is potentially one of the main mechanisms through which light/ day length information is used to control or alter the clock. [The central oscillator in turn regulates PIF expression. Postdusk, TOC1 peaks in expression, allowing direct interactions between TOC1 and PIF3, which results in the TOC1 gating of PIF induced growth, until TOC1 levels decrease predawn \(Soy *et al.*, 2016\).](#)

Formatted: Font: Italic

PIF4 and 5 show rhythmic expression with a diurnal peak at dawn (Nomoto *et al.*, 2012). Expression of *PIF4* and 5 is controlled by the evening complex, comprising of ELF3, ELF4 and LUX (Herrero *et al.*, 2012), which binds to the promoter region of *PIF4* and 5 to inhibit transcription at dusk (Nusinow *et al.*, 2011). PIF protein stably accumulates overnight due to the lack of phytochrome induced degradation to reach their maximum level at dawn (Shin *et al.*, 2013). As PIFs are growth-promoting factors (Shin *et al.*, 2013), this leads to the highest growth rate occurring at the end of the night phase.


PIFs may also input information from other environmental cues to the clock such as temperature (McClung & Davis, 2010; Raschke *et al.* 2015). It was shown that *PIF4* expression also increases in response to temperature increases (Shin *et al.*, 2013). As dawn induces a temperature increase, alongside the return of daylight it would perhaps be advantageous to a plant to be able to input both of these environmental cues into the clock at the same time.

PIF3 like (PILs) are also basic helix loop helix transcription factors with large overlaps in function to PIFs, but have been associated with shade avoidance (Li *et al.*, 2014). This overlap in function has led to some ambiguous nomenclature as PIL5 is also referred to as PIF1 and likewise PIL6 as PIF5 (Li *et al.*, 2014). PIL1 has a distinct function and was shown to work with TOC1 to restrict growth at specific times of day (Salter *et al.*, 2003). PIL1 accumulates rapidly within the first hour of shade cover acting as part of a rapid signalling pathway to stop growth (Li *et al.*, 2014), a secondary longer lasting shade response is then mediated by HFR and phytochrome rapidly regulated (PAR1/2) (Galstyan *et al.*, 2011). The exact mechanism through which PIL1 halts growth in shade is not known but a number of hypothesis were presented in (Li *et al.*, 2014). It was suggested that as PIF1 has a binding site for phyB it is possible that in shade PIL1 may outcompete PIF for DNA binding sites on the Pfr form of phyB, therefore reducing the growth promoting function of PIF5 (Li *et al.*, 2014). Alternatively, PIL1 may work via a PIF independent mechanism on components of downstream pathways; however this is yet to be tested.

HFR

Long hypocotyl in far red 1 (HFR1) is a basic helix loop helix transcription factor involved in phytochrome-mediated signalling (Fairchild *et al.*, 2000), and photomorphogenesis (Yang *et al.*, 2005). HFR is unstable in darkness and accumulates in the light, this accumulation is due to light preventing COP1 mediated degradation of HFR (Yang *et al.*, 2005). HFR is not able to bind phyA or B directly, instead HFR binds PIF3 forming potentially a heterodimer of PIF3/HFR which can then bind to the Pfr form of phyA/B. This is also highlighted by the fact that HFR is more abundantly found in far red light (Fairchild *et al.*, 2000). Mutants deficient in HFR had defective phyA responses, such as hypocotyl elongation, and induction of chlorophyll A binding protein (CAB) (Fankhauser & Chory, 2000). HFR is also thought to have a blue-light response (Duek & Fankhauser, 2003) through CRY1 (Yang *et al.*, 2005), but the exact mechanism through which this occurs is not known.

For Review Only



Intersection of the clock components ELF3- and ELF4 to light and clock signalling: major integrators of light to the clock

EARLY FLOWERING 3 (ELF3) was first identified as a negative regulator of flowering time. In addition to the observation that *elf3* mutant was shown to be early flowering, large circadian defects were identified (Hicks *et al.*, 1996, Roden *et al.*, 2002; Undurraga *et al.*, 2012). *elf3* mutants are defective in gating of red-light perception to the clock. Cloning of ELF3 allowed for interactors to be detected, phyB was revealed to be a factor that associated to the N-terminus of ELF3 (Liu *et al.*, 2001). ELF3 was identified as playing a role in light signalling, in 12hours light 12 hours dark entrainment, ELF3 accumulates in the nucleus just before darkness (ZT12) (Liu *et al.*, 2001). Increasing day length, increases the nuclear accumulation of ELF3, increased darkness causes accumulation of ELF3 to decrease to an undetectable level (Liu *et al.*, 2001) showing a direct relationship between light and ELF3.

Phase response curves are made by measuring circadian period and phase during light pulses, at times across subjective day and night. Phase response curves for wild-type Arabidopsis in both red and blue light show the greatest effect and subsequent clock resetting to be caused by a light pulse during the subjective night. *ELF3* overexpression lines showed a much more gradual change in phase response with the same light pulses (Covington *et al.*, 2001). In white light, *ELF3* overexpression causes a period lengthening effect in a light-dependent manner. In darkness, the oscillator pace is not altered (Covington *et al.*, 2001). In *elf3* mutants, phase response light pulses showed a much greater effect than the wild type, suggesting that clock resetting is light dependent (Covington *et al.*, 2001), involving ELF3 in oscillator resetting by repressing the light input to the clock (Bujdosó & Davis, 2013). However, the exact mechanism through which this occurs is unknown.

elf3 and *phyB* mutants were found to have similar phenotypic traits, such as hypocotyl elongation in red light and constitutive shade avoidance (Devlin *et al.*, 1999). *elf3* mutants are also defective in their response to blue and/ or red light with stronger effects showing in red light. ELF3 grown in darkness is rhythmic with a long-period phenotype, rhythmic in light/dark entrainment cycles, but arrhythmic in free run light conditions (Hicks *et al.*, 1996). *elf3* plants are also unable to inhibit hypocotyl elongation under light (Zagotta *et al.*, 1996). The combined phenotypic characteristics of the *elf3* mutants suggest that ELF3 plays a role in light perception and signalling, particularly in red light due to the interaction between ELF3 and phyB (Reed *et al.*, 2000).

EARLY FLOWERING 4 (ELF4) was the first clock component interpreted as being required for the clock to cycle and it was revealed to be a component of normal light perception. Genetic loss of *ELF4* resulted in plants that were markedly attenuated in the ability of a red-light pulse to generate *CCA1* and *LHY* rhythms in etiolated plants. This was concluded to be due to ELF4 being required for the phytochrome-mediated light induction of *CCA1* and *LHY* expression (Kikis *et al.*, 2005). Consistent with this, it was shown that *elf4* mutants were hypermorphic and hypomorphic to red-light cues, dependent on the assay. Notably, red-light mediated induction of *CAB2* expression was elevated in *elf4* (McWatters *et al.*, 2007). This revealed that ELF4 contributes to so-called circadian gating of light responsiveness (negative photomorphogenesis), light-regulation of *PIF4/5* expression and the suppression of growth (positive photomorphogenesis) (Nozue *et al.*, 2007).

The presence of ELF4 in the nucleus increases the accumulation of ELF3 (Herrero *et al.*, 2012). ELF4 may function as a nuclear anchor for ELF3, but does not affect the nuclear localisation of LUX, the third component of the evening complex (Herrero *et al.*, 2012). Nuclear import of phyB is light dependent (Kircher *et al.*, 1999, Sakamoto & Nagatani, 1996), and it was shown that phyB does not import ELF3 into the nucleus (Bujdosó & Davis, 2013). It is possible that ELF3 plays a role in the nuclear import of phyB, as the N-terminus of ELF3 interacts with the C-terminal end of both the Pr and Pfr forms of PHYB (Liu *et al.*, 2001). However, ELF3 and phyB have opposite roles in controlling circadian oscillations (Herrero *et al.*, 2012). ELF3 needs the association with ELF4 to maintain circadian oscillations, counteracting the COP1 and phyB mediated repression of ELF3 (Herrero *et al.*, 2012). ELF3 is also part of the blue-light signalling pathway, through its interaction with COP1. How

phyB and ELF4 coordinate the action and localisation of ELF3 seems critical for the cooperative intersection of light perception and circadian clock function.

It was originally unclear what overall effect ELF3 has on other clock genes as the *elf3* loss of function mutation causes arrhythmicity. A reduction-of-function mutation in *elf3-12* provided a way to explore this, as the hypomorphic *elf3-12* allele is able to maintain rhythmicity (Kolmos *et al.*, 2011). This showed *elf3-12* to be light dependent but with a defective phase resetting mechanism. This *elf3-12* mutant allowed the position of ELF3 within the clock to be derived as key to the regulation of *PRR9* expression (Kolmos & Davis, 2007, Kolmos *et al.*, 2011). It is known that ELF3 associates to the promoter of *PRR9* to mediate its repression. In addition to clock-regulated *PRR9* transcription, *PRR9* expression is also light regulated. Furthermore, *prp9* mutants display photomorphogenic phenotypes (Nakamichi *et al.*, 2005). This highlights the role of ELF3 in reciprocally linking light signalling to clock function.

Overexpression of *PHYA* in an *elf3-12* background showed that the overexpressed *PHYA* has an additive effect with the *elf3-12* mutation to give a further shortened period in red light. In a range of light conditions the *elf3-12 PHYA*-overexpression lines had an altered phase however in darkness there was no change suggesting that light has an epistatic effect on *PHYA* (Kolmos *et al.*, 2011). Overexpression of *PHYB* in the same *elf3-12* background gave the same period shortening as the *PHYA*-overexpression lines; however the *PHYB-elf3-12* lines had a phase more closely linked to *PHYB* overexpression in a wild-type background (Kolmos *et al.*, 2011). This result suggests that *PHYB* functions upstream of *elf3-12* in light signalling and is able to suppress the effects of *elf3-12* (Bujdoso & Davis, 2013). Together it appears that some, but not all, inputs of phytochromes to the clock depend on ELF3.

Duration and quality of light

In regular light-dark cycles Arabidopsis has a circadian period of approximately 24 hours in light, whereas in darkness it has a period of 30 to 36 hours (Millar *et al.*, 1995). Light therefore makes the clock run faster, the absence of light cues causes the clock to slow, this is in keeping with Aschoff's rule (Aschoff, 1979). The range of photoreceptors present in Arabidopsis allow a range of fluence rates to be detected, ensuring the phase of the circadian oscillator is synchronised with environmental cues (Somers *et al.*, 1998a), at both dawn and dusk (Devlin & Kay, 2000a). Removal or even partial reduction of blue and red photoreceptors causes the clock to run slower (Millar *et al.*, 1995), suggesting that the effects of light intensity on the speed of the clock is limited by the number of photoreceptors present. This would also suggest that it would not be possible to increase the speed of the clock with higher light intensities beyond the maximum speed obtainable by that number of photoreceptors.

Input of duration and quality of light are important in synchronising processes such as flowering time and development (Weston *et al.*, 2000). Preceding photoperiod was shown to alter the subsequent speed of the clock (Darrah *et al.*, 2006; Boikoglou *et al.* 2011). Interestingly here is the long known role of ELF3 in processing light information to the clock (Hicks *et al.* 1996), and how this coordinates the capacity for a plant to perceive daily boundaries present in a day night cycle (McWatters *et al.* 2000). Recent work has revealed that extensive allelic variation at *ELF3* contribute to alterations in photoperiodic control and this is associated to alterations in encoded nuclear abundance and *in vivo* turn over diurnal time (Undurraga *et al.* 2012, Anwer *et al.* 2014). Combined with temperature variation over the day, light duration gives information on the time of year or season and therefore a warning of the growth conditions to follow. It is clear that allelic variation exists in Arabidopsis to change the output of such varying entrainment processes (Darrah *et al.*, 2006, Boikoglou *et al.* 2011; Anwer *et al.* 2014).

Plants are able to adapt to changes in light intensity, such as consistently low light intensity, by re-arranging photosynthetic machinery to be more efficient at light harvesting (Weston *et al.*, 2000). Blue light plays a major role in this. As such it would be assumed that cryptochromes and *PHYA/B* play a role in directing the timing of light capture. A connection hub for this, COP1 has been

identified as a signalling intermediate between these two processes (Walters *et al.*, 1999), however links between photoreceptor function and organisation of the photosynthetic apparatus await further investigation (Walters *et al.*, 1999, Weston *et al.*, 2000). Finally light intensity is also detected by the plastid sensing blue light, causing structural changes and elongation of the palisade to absorb more light, this process responds a lot more to blue light than red (Weston *et al.*, 2000). It could be hypothesised that this plastid information is used as a nuclear clock and is synchronised with the phyA and phyB red light input to the central oscillator by COP1.

Discussion

Multiple photoreceptors are essential components of light input to the clock. In this way they play a central role in the light input to the clock. Not only with the reversible, light-mediated reactions for maximal efficiency in light or dimmer light / shade, they also directly input light into the central oscillator through clock associated factors, such as ELF3/4, COP1, ZTL, PIFs, PILs and HFR. Transcriptional regulation and post-translational processes are all part of this complex web of interconnections between light-perception and clock function. Overall light input to the clock forms complex feedback systems that generated harmonised regulatory pathways, the mechanisms from light perception to clock function, and back again, awaits clear discoveries.

Many plant growth chambers are produced with red, blue and far-red LED panels, but are these the optimal light regimes to measure plant gene expression under? Furthermore, most chambers have a lights-on or off function which does not represent the graded changes in intensity that would occur with a plant growing under natural sunlight (or in a greenhouse), with the gradual appearance and disappearance of light at sunrise and sunset. As white light comprises a combination of different wavelengths of light simultaneously, it may be possible that the pathways for different colours of light interact more than has been found so far. Overlapping functions have been found between blue and red light and these are the most commonly used light wavelengths for plant LEDs. It is possible that there are essential components missing in just red and blue, although difficult to isolate in a complex web of circadian gene expression, may only be present in white light or in other light combinations not yet tested.

Light is essential for plant growth and it is therefore important to understand how plants process the daily light cues they receive. Further understanding of how each light wavelength is detected and the information fed into the central oscillator from each sensor, could potentially have a large impact on plant and ultimately crop growth. With global changes in climate, knowledge on essential lighting requirements and how this impacts on overall plant health could be used to optimise crop productivity. Indoor farming techniques using LEDs as a light source could be optimised for maximal yield and growth speed. As permafrost regions recede, and suitable growth land becomes available. A detailed understanding of photoperiodicity and how it impacts fitness will help with the challenges created by growing crops in shorter growth seasons and longer daylight hours.

Formatted: Indent: First line: 0.5"

Figure legends

Figure 1. Schematic diagram of feedback loops in the Arabidopsis central oscillator.

Figure 2. Effects of light intensity on circadian period. A). Circadian gene expression is rhythmic in constant light but in the absence of exogenous sucrose, rapidly becomes arrhythmic in prolonged darkness. B). Following Aschoff's rule, high intensity light causes the clock to run faster; lower light intensities cause the clock to slow. C). High light intensities result in period shortening; lower light intensities result in period lengthening.

Figure 3. Circadian photoreceptors and their potential interactions to input light to the central oscillator.

- Formatted: Font: Bold
- Formatted: Font: Bold
- Formatted: Font: Bold
- Formatted: Font: Bold, Font color: Auto
- Formatted: Normal, Left
- Formatted: Font: 11 pt, Font color: Auto
- Formatted: Font: 11 pt, Not Bold, Font color: Auto
- Formatted: Font: 11 pt
- Formatted: Font: 11 pt, Font color: Auto
- Formatted: Font: 11 pt, Not Bold, Font color: Auto
- Formatted: Font: 11 pt

References

- Adams S., Manfield I., Stockley P. & Carré I.A. (2015) Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. *PLOS ONE*, **10**, e0143943.
- Ahmad M. & Cashmore A.R. (1993) HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature*, **366**, 162-166.
- Ahmad M., Jarillo J.A. & Cashmore A.R. (1998a) Chimeric Proteins between cry1 and cry2 Arabidopsis Blue Light Photoreceptors Indicate Overlapping Functions and Varying Protein Stability. *The Plant Cell*, **10**, 197-207.
- Ahmad M., Jarillo J.A., Smirnova O. & Cashmore A.R. (1998b) The CRY1 Blue Light Photoreceptor of Arabidopsis Interacts with Phytochrome A In Vitro. *Molecular Cell*, **1**, 939-948.
- Alabadí D., Oyama T., Yanovsky M.J., Harmon F.G., Más P. & Kay S.A. (2001) Reciprocal Regulation Between TOC1 and LHY/CCA1 Within the Arabidopsis Circadian Clock. *Science*, **293**, 880-883.
- Anderson S.L., Somers D.E., Millar A.J., Hanson K., Chory J. & Kay S.A. (1997) Attenuation of phytochrome A and B signaling pathways by the Arabidopsis circadian clock. *The Plant Cell*, **9**, 1727-1743.
- Anwer M.U., Boikoglou E., Herrero E., Hallstein M., Davis A.M., Velikkakam James G., Nagy F. & Davis S.J. (2014) Natural variation reveals that intracellular distribution of ELF3 protein is associated with function in the circadian clock. *eLife*, e02206.
- Aschoff J. (1979) Circadian Rhythms: Influences of Internal and External Factors on the Period Measured in Constant Conditions. *Zeitschrift für Tierpsychologie*, **49**, 225-249.
- Aukerman M.J., Hirschfeld M., Wester L., Weaver M., Clack T., Amasino R.M. & Sharrock R.A. (1997) A deletion in the PHYD gene of the Arabidopsis Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *The Plant Cell*, **9**, 1317-1326.
- Banerjee R., Schleicher E., Meier S., Viana R.M., Pokorny R., Ahmad M., Bittl R. & Batschauer A. (2007) The Signaling State of Arabidopsis Cryptochrome 2 Contains Flavin Semiquinone. *Journal of Biological Chemistry*, **282**, 14916-14922.
- Baudry A., Ito S., Song Y.H., Strait A.A., Kiba T., Lu S., Henriques R., Pruneda-Paz J.L., Chua N.H., Tobin E.M., Kay S.A., Imaizumi T. (2010) F-box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control Arabidopsis clock progression. *The Plant Cell*, **22**, 606-622.
- Bläsing O.E., Gibon Y., Günther M., Höhne M., Morcuende R., Osuna D., Thimm O., Usadel B., Scheible W.-R. & Stitt M. (2005) Sugars and Circadian Regulation Make Major Contributions to the Global Regulation of Diurnal Gene Expression in Arabidopsis. *The Plant Cell*, **17**, 3257-3281.
- Bognár L.K., Hall A., Ádám É., Thain S.C., Nagy F. & Millar A.J. (1999) The circadian clock controls the expression pattern of the circadian input photoreceptor, phytochrome B. *Proceedings of the National Academy of Sciences of the United States of America*, **96**, 14652-14657.
- Boikoglou E., Ma Z., von Korff M., Davis A.M., Nagy F. & Davis S.J. (2011) Environmental memory from a circadian oscillator: the Arabidopsis thaliana clock differentially integrates perception of photic vs. thermal entrainment. *Genetics*, **189**, 655-664.
- Bouly J.P., Schleicher E., Dionisio-Sese M., Vandenbussche F., Van Der Straeten D., Bakrim N., Meier S., Batschauer A., Galland P., Bittl R. & Ahmad M. (2007) Cryptochrome Blue Light Photoreceptors Are Activated through Interconversion of Flavin Redox States. *Journal of Biological Chemistry*, **282**, 9383-9391.
- Brown B.A. & Jenkins G.I. (2008) UV-B Signaling Pathways with Different Fluence-Rate Response Profiles Are Distinguished in Mature Arabidopsis Leaf Tissue by Requirement for UVR8, HY5, and HYH. *Plant Physiology*, **146**, 576-588.
- Bujdoso N. & Davis S. (2013) Mathematical modeling of an oscillating gene circuit to unravel the circadian clock network of Arabidopsis thaliana. *Frontiers in Plant Science*, **4**.

Formatted: Font: Calibri, 11 pt, Font color: Auto, English (U.S.), Do not check spelling or grammar

Formatted: English (U.S.), Do not check spelling or grammar

- Carre I.A. & Kay S.A. (1995) Multiple DNA-Protein Complexes at a Circadian-Regulated Promoter Element. *The Plant Cell*, **7**, 2039-2051.
- Cashmore A.R., Jarillo J.A., Wu Y.J. & Liu D. (1999) Cryptochromes: Blue Light Receptors for Plants and Animals. *Science*, **284**, 760-765.
- Cha J.Y., Kim J., Kim T.S., Zeng Q., Wang L., Lee S.Y., Kim W.Y. & Somers D.E. (2017) GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE in the Arabidopsis circadian clock. *Nature Communications*, **8**, 3.
- Chow B.Y., Sanchez S.E., Breton G., Pruneda-Paz J.L., Krogan N.T. & Kay S.A. (2014) Transcriptional Regulation of LUX by CBF1 Mediates Cold Input to the Circadian Clock in Arabidopsis. *Current Biology*, **24**, 1518-1524.
- Clough R.C. & Vierstra R.D. (1997) Phytochrome degradation. *Plant, Cell & Environment*, **20**, 713-721.
- Covington M.F., Panda S., Liu X.L., Strayer C.A., Wagner D.R. & Kay S.A. (2001) ELF3 Modulates Resetting of the Circadian Clock in Arabidopsis. *The Plant Cell*, **13**, 1305-1316.
- Darrah C., Taylor B.L., Edwards K.D., Brown P.E., Hall A. & McWatters H.G. (2006) Analysis of Phase of LUCIFERASE Expression Reveals Novel Circadian Quantitative Trait Loci in Arabidopsis. *Plant Physiology*, **140**, 1464-1474.
- Davis S.J. (2002) Photoperiodism: The Coincidental Perception of the Season. *Current Biology*, **12**, R841-R843.
- Delker C., Sonntag L., James G.V., Janitzka P., Ibañez C., Ziermann H., Peterson T., Denk K., Mull S., Ziegler J., Davis S.J., Schneeberger K. & Quint M. (2014) The DET1-COP1-HY5 pathway constitutes a multipurpose signaling module regulating plant photomorphogenesis and thermomorphogenesis. *Cell Rep.* **24**:1983-1989.
- Devlin P.F. & Kay S.A. (2000a) Cryptochromes are required for phytochrome signaling to the circadian clock but not for rhythmicity. *Plant Cell*, **12**, 2499-2510.
- Devlin P.F. & Kay S.A. (2000b) Flower arranging in Arabidopsis. *Science*, **288**, 1600-1602.
- Devlin P.F., Patel S.R. & Whitelam G.C. (1998) Phytochrome E Influences Internode Elongation and Flowering Time in Arabidopsis. *The Plant Cell*, **10**, 1479-1487.
- Devlin P.F., Robson P.R.H., Patel S.R., Goosey L., Sharrock R.A. & Whitelam G.C. (1999) Phytochrome D Acts in the Shade-Avoidance Syndrome in Arabidopsis by Controlling Elongation Growth and Flowering Time. *Plant Physiology*, **119**, 909-916.
- Dodd A.N., Salathia N., Hall A., Kevei E., Toth R., Nagy F., Hibberd J.M., Millar A.J. & Webb A.A.R. (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. *Science*, **309**, 630-633.
- Duek P.D. & Fankhauser C. (2003) HFR1, a putative bHLH transcription factor, mediates both phytochrome A and cryptochrome signalling. *The Plant Journal*, **34**, 827-836.
- Edwards K.D., Guerineau F., Devlin P.F. & Millar A.J. (2015) Low temperature effects of *PHYTOCHROME C* on the circadian clock in Arabidopsis suggest that *PHYC* underlies natural variation in biological timing. *bioRxiv*.
- Fairchild C.D., Schumaker M.A. & Quail P.H. (2000) HFR1 encodes an atypical bHLH protein that acts in phytochrome A signal transduction. *Genes & Development*, **14**, 2377-2391.
- Fankhauser C. & Chory J. (2000) RSF1, an Arabidopsis Locus Implicated in Phytochrome A Signaling. *Plant Physiology*, **124**, 39-46.
- Favory J.J., Stec A., Gruber H., Rizzini L., Oravecz A., Funk M., Albert A., Cloix C., Jenkins G.I., Oakeley E.J., Seidlitz H.K., Nagy F. & Ulm R. (2009) Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in Arabidopsis. *The EMBO Journal*, **28**, 591-601.
- Federer C.A. & Tanner C.B. (1966) Spectral Distribution of Light in the Forest. *Ecology*, **47**, 555-560.
- Fehér B., Kozma-Bognár L., Kevei É., Hajdu A., Binkert M., Davis S.J., Schäfer E., Ulm R. & Nagy F. (2011) Functional interaction of the circadian clock and UV RESISTANCE LOCUS 8-controlled UV-B signaling pathways in Arabidopsis thaliana. *The Plant Journal*, **67**, 37-48.

- Findlay K.M.W. & Jenkins G.I. (2016) Regulation of UVR8 photoreceptor dimer/monomer photo-equilibrium in Arabidopsis plants grown under photoperiodic conditions. *Plant, Cell & Environment*, **39**, 1706-1714.
- Folta K.M. (2004) Green Light Stimulates Early Stem Elongation, Antagonizing Light-Mediated Growth Inhibition. *Plant Physiology*, **135**, 1407-1416.
- Folta K.M. & Maruhnich S.A. (2007) Green light: a signal to slow down or stop. *Journal of Experimental Botany*, **58**, 3099-3111.
- Fornara F., de Montaigu A., Sanchez-Villarreal A., Takahashi Y., van Themaat E.V.L., Huettel B., Davis S.J. & Coupland G. (2015) The GI-CDF module of Arabidopsis affects freezing tolerance and growth as well as flowering. *Plant Journal*, **81**, 695-706.
- Franklin K.A., Davis S.J., Stoddart W.M., Vierstra R.D. & Whitelam G.C. (2003) Mutant Analyses Define Multiple Roles for Phytochrome C in Arabidopsis Photomorphogenesis. *The Plant Cell*, **15**, 1981-1989.
- Franklin K.A. & Quail P.H. (2010) Phytochrome functions in Arabidopsis development. *Journal of Experimental Botany*, **61**, 11-24.
- Fujiwara S., Wang L., Han L., Suh S. S., Salomé P. A., McClung C. R., Somers D. E. (2008) Post-translational regulation of the Arabidopsis circadian clock through selective proteolysis and phosphorylation of pseudo-response regulator proteins. *Journal of Biological Chemistry*, **283**, 23073-23083.
- Galstyan A., Cifuentes-Esquivel N., Bou-Torrent J. & Martinez-Garcia J.F. (2011) The shade avoidance syndrome in Arabidopsis: a fundamental role for atypical basic helix-loop-helix proteins as transcriptional cofactors. *The Plant Journal*, **66**, 258-267.
- Gendron J.M., Pruneda-Paz J.L., Doherty C.J., Gross A.M., Kang S.E. & Kay S.A. (2012) Arabidopsis circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proceedings of the National Academy of Sciences*, **109**, 3167-3172.
- Goto N., Kumagai T. & Koornneef M. (1991) Flowering responses to light-breaks in photomorphogenic mutants of Arabidopsis thaliana, a long-day plant. *Physiologia Plantarum*, **83**, 209-215.
- Guerriero M.L., Pokhilko A., Fernández A.P., Halliday K.J., Millar A.J. & Hillston J. (2012) Stochastic properties of the plant circadian clock. *Journal of the Royal Society Interface*, **9**, 744-756.
- Guo H., Yang H., Mockler T.C. & Lin C. (1998) Regulation of Flowering Time by Arabidopsis Photoreceptors. *Science*, **279**, 1360-1363.
- Habte E., Muller L.M., Shtaya M., Davis S.J. & von Korff M. (2014) Osmotic stress at the barley root affects expression of circadian clock genes in the shoot. *Plant Cell and Environment*, **37**, 1321-1337.
- Hanano S., Stracke R., Jakoby M., Merkle T., Domagalska M.A., Weisshaar B. & Davis S.J. (2008) A systematic survey in Arabidopsis thaliana of transcription factors that modulate circadian parameters. *BMC Genomics*, **9**, 182.
- Harmer S.L. (2009) The Circadian System in Higher Plants. *Annual Review of Plant Biology*, **60**, 357-377.
- Harmer S.L., Hogenesch J.B., Straume M., Chang H.S., Han B., Zhu T., Wang X., Kreps J.A. & Kay S.A. (2000) Orchestrated Transcription of Key Pathways in Arabidopsis by the Circadian Clock. *Science*, **290**, 2110-2113.
- Haydon M.J., Mielczarek O., Robertson F.C., Hubbard K.E. & Webb A.A.R. (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. *Nature*, **502**, 689-692.
- Haydon M.J. & Webb A.A. (2016) Assessing the Impact of Photosynthetic Sugars on the Arabidopsis Circadian Clock. *Environmental Responses in Plants: Methods and Protocols*, 133-140.
- Hayes S., Sharma A., Fraser D.P., Trevisan M., Cragg-Barber C.K., Tavridou E., Fankhauser C., Jenkins G.I. & Franklin K.A. (2017) UV-B Perceived by the UVR8 Photoreceptor Inhibits Plant Thermomorphogenesis. *Current Biology*, **27**, 120-127.

Formatted: English (U.S.), Do not check spelling or grammar

- Hazen S.P., Schultz T.F., Pruneda-Paz J.L., Borevitz J.O., Ecker J.R. & Kay S.A. (2005) LUX ARRHYTHMO encodes a Myb domain protein essential for circadian rhythms. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 10387-10392.
- Heijde M. & Ulm R. (2012) UV-B photoreceptor-mediated signalling in plants. *Trends in Plant Science*, **17**, 230-237.
- Helfer A., Nusinow D.A., Chow B.Y., Gehrke A.R., Bulyk M.L. & Kay S.A. (2011) LUX ARRHYTHMO Encodes a Nighttime Repressor of Circadian Gene Expression in the Arabidopsis Core Clock. *Current Biology*, **21**, 126-133.
- Herrero E., Kolmos E., Bujdoso N., Yuan Y., Wang M., Berns M.C., Uhlworm H., Coupland G., Saini R., Jaskolski M., Webb A., Gonçalves J. & Davis S.J. (2012) EARLY FLOWERING4 Recruitment of EARLY FLOWERING3 in the Nucleus Sustains the Arabidopsis Circadian Clock. *The Plant Cell*, **24**, 428-443.
- Hicks K.A., Millar A.J., Carré I.A., Somers D.E., Straume M., Meeks-Wagner D.R. & Kay S.A. (1996) Conditional Circadian Dysfunction of the Arabidopsis early-flowering 3 Mutant. *Science*, **274**, 790-792.
- Hoecker U. & Quail P.H. (2001) The Phytochrome A-specific Signaling Intermediate SPA1 Interacts Directly with COP1, a Constitutive Repressor of Light Signaling in Arabidopsis. *Journal of Biological Chemistry*, **276**, 38173-38178.
- Hoecker U., Tepperman J.M. & Quail P.H. (1999) SPA1, a WD-Repeat Protein Specific to Phytochrome A Signal Transduction. *Science*, **284**, 496.
- Hu W., Franklin K.A., Sharrock R.A., Jones M.A., Harmer S.L. & Lagarias J.C. (2013) Unanticipated regulatory roles for Arabidopsis phytochromes revealed by null mutant analysis. *Proceedings of the National Academy of Sciences*, **110**, 1542-1547.
- Huang H., Alvarez S., Bindbeutel R., Shen Z., Naldrett M.J., Evans B.S., Briggs S.P., Hicks L.M., Kay S.A. & Nusinow D.A. (2016) Identification of Evening Complex Associated Proteins in Arabidopsis by Affinity Purification and Mass Spectrometry. *Molecular & Cellular Proteomics*, **15**, 201-217.
- Huang X., Ouyang X. & Deng X.W. (2014) Beyond repression of photomorphogenesis: role switching of COP/DET/FUS in light signaling. *Current Opinion in Plant Biology*, **21**, 96-103.
- Jang S., Marchal V., Panigrahi K.C.S., Wenkel S., Soppe W., Deng X.-W., Valverde F. & Coupland G. (2008) Arabidopsis COP1 shapes the temporal pattern of CO accumulation conferring a photoperiodic flowering response. *The EMBO Journal*, **27**, 1277-1288.
- Jansen M.A.K., Gaba V. & Greenberg B.M. (1998) Higher plants and UV-B radiation: balancing damage, repair and acclimation. *Trends in Plant Science*, **3**, 131-135.
- Jarillo J.A., Capel J., Tang R.H., Yang H.Q., Alonso J.M., Ecker J.R. & Cashmore A.R. (2001) An Arabidopsis circadian clock component interacts with both CRY1 and phyB. *Nature*, **410**, 487-490.
- Jiménez-Gómez J.M., Wallace A.D. & Maloof J.N. (2010) Network analysis identifies ELF3 as a QTL for the shade avoidance response in Arabidopsis. *PLoS Genet*, **6**, e1001100.
- Johnson C.H., Knight M.R., Kondo T., Masson P., Sedbrook J., Haley A. & Trewavas A. (1995) Circadian oscillations of cytosolic and chloroplastic free calcium in plants. *Science*, **269**, 1863.
- Kamioka M., Takao S., Suzuki T., Taki K., Higashiyama T., Kinoshita T. & Nakamichi N. (2016) Direct Repression of Evening Genes by CIRCADIEN CLOCK-ASSOCIATED1 in the Arabidopsis Circadian Clock. *The Plant Cell*, **28**, 696-711.
- Kevei É., Gyula P., Hall A., Kozma-Bognár L., Kim W.-Y., Eriksson M.E., Tóth R., Hanano S., Fehér B., Southern M.M., Bastow R.M., Viczián A., Hibberd V., Davis S.J., Somers D.E., Nagy F. & Millar A.J. (2006) Forward Genetic Analysis of the Circadian Clock Separates the Multiple Functions of ZEITLUPE. *Plant Physiology*, **140**, 933-945.
- Kiba T., Henriques R., Hitoshi S., Chua, N.H. (2007) Targeted Degradation of PSEUDO-RESPONSE REGULATOR5 by an SCF^{TRT} Complex Regulates Clock Function and Photomorphogenesis in Arabidopsis thaliana. *The Plant Cell*, **19** (8) 2516-2530.

Formatted: Font: Calibri, 11 pt, Font color: Auto, English (U.S.), Do not check spelling or grammar

Formatted: English (U.S.), Do not check spelling or grammar

- Kikis E.A., Khanna R. & Quail P.H. (2005) ELF4 is a phytochrome-regulated component of a negative-feedback loop involving the central oscillator components CCA1 and LHY. *The Plant Journal*, **44**, 300-313.
- Kim T.s., Kim W.Y., Fujiwara S., Kim J., Cha J.Y., Park J.H., Lee S.Y. & Somers D.E. (2011) HSP90 functions in the circadian clock through stabilization of the client F-box protein ZEITLUPE. *Proceedings of the National Academy of Sciences*, **108**, 16843-16848.
- Kim W.Y., Fujiwara S., Suh S.S., Kim J., Kim Y., Han L., David K., Putterill J., Nam H.G. & Somers D.E. (2007) ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature*, **449**, 356-360.
- Kim W.Y., Hicks K.A. & Somers D.E. (2005) Independent Roles for EARLY FLOWERING 3 and ZEITLUPE in the Control of Circadian Timing, Hypocotyl Length, and Flowering Time. *Plant Physiology*, **139**, 1557-1569.
- Kircher S., Kozma-Bognar L., Kim L., Adam E., Harter K., Schäfer E. & Nagy F. (1999) Light Quality-Dependent Nuclear Import of the Plant Photoreceptors Phytochrome A and B. *The Plant Cell*, **11**, 1445-1456.
- Kolmos E. & Davis S.J. (2007) ELF4 as a Central Gene in the Circadian Clock. *Plant Signaling & Behavior*, **2**, 370-372.
- Kolmos E., Herrero E., Bujdoso N., Millar A.J., Toth R., Gyula P., Nagy F. & Davis S.J. (2011) A Reduced-Function Allele Reveals That EARLY FLOWERING3 Repressive Action on the Circadian Clock Is Modulated by Phytochrome Signals in Arabidopsis. *The Plant Cell*, **23**, 3230-3246.
- Koornneef M., Rolff E. & Spruit C.J.P. (1980) Genetic Control of Light-inhibited Hypocotyl Elongation in Arabidopsis thaliana (L.) Heynh. *Zeitschrift für Pflanzenphysiologie*, **100**, 147-160.
- Laubinger S., Fittinghoff K. & Hoecker U. (2004) The SPA Quartet: A Family of WD-Repeat Proteins with a Central Role in Suppression of Photomorphogenesis in Arabidopsis. *The Plant Cell*, **16**, 2293-2306.
- Leivar P., Monte E., Cohn M.M. & Quail P.H. (2012) Phytochrome Signaling in Green Arabidopsis Seedlings: Impact Assessment of a Mutually Negative phyB-PIF Feedback Loop. *Molecular Plant*, **5**, 734-749.
- Li L., Zhang Q., Pedmale U.V., Nito K., Fu W., Lin L., Hazen S.P. & Chory J. (2014) PIL1 Participates in a Negative Feedback Loop that Regulates Its Own Gene Expression in Response to Shade. *Molecular Plant*, **7**, 1582-1585.
- Lian H.L., He S.B., Zhang Y.C., Zhu D.M., Zhang J.Y., Jia K.P., Sun S.X., Li L. & Yang H.Q. (2011) Blue-light-dependent interaction of cryptochrome 1 with SPA1 defines a dynamic signaling mechanism. *Genes & Development*, **25**, 1023-1028.
- Lin C., Ahmad M. & Cashmore A.R. (1996) Arabidopsis cryptochrome 1 is a soluble protein mediating blue light-dependent regulation of plant growth and development. *The Plant Journal*, **10**, 893-902.
- Lin C., Yang H., Guo H., Mockler T., Chen J. & Cashmore A.R. (1998) Enhancement of blue-light sensitivity of Arabidopsis seedlings by a blue light receptor cryptochrome 2. *Proceedings of the National Academy of Sciences*, **95**, 2686-2690.
- Liu H., Yu X., Li K., Klejnot J., Yang H., Lisiero D. & Lin C. (2008) Photoexcited CRY2 Interacts with CIB1 to Regulate Transcription and Floral Initiation in Arabidopsis. *Science*, **322**, 1535-1539.
- Liu X.L., Covington M.F., Fankhauser C., Chory J. & Wagner D.R. (2001) ELF3 Encodes a Circadian Clock-Regulated Nuclear Protein That Functions in an Arabidopsis PHYB Signal Transduction Pathway. *The Plant Cell*, **13**, 1293-1304.
- Locke J.C.W., Kozma-Bognár L., Gould P.D., Fehér B., Kevei É., Nagy F., Turner M.S., Hall A. & Millar A.J. (2006) Experimental validation of a predicted feedback loop in the multi-oscillator clock of Arabidopsis thaliana. *Molecular Systems Biology*, **2**.

- Lorrain S., Allen T., Duek P.D., Whitelam G.C. & Fankhauser C. (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *The Plant Journal*, **53**, 312-323.
- Martínez-García J.F., Huq E. & Quail P.H. (2000) Direct Targeting of Light Signals to a Promoter Element-Bound Transcription Factor. *Science*, **288**, 859-863.
- Más P., Alabadí D., Yanovsky M.J., Oyama T. & Kay S.A. (2003) Dual Role of TOC1 in the Control of Circadian and Photomorphogenic Responses in Arabidopsis. *The Plant Cell*, **15**, 223-236.
- Mas P., Devlin P.F., Panda S. & Kay S.A. (2000) Functional interaction of phytochrome B and cryptochrome 2. *Nature*, **408**, 207-211.
- Mas P., Kim W.Y., Somers D.E. & Kay S.A. (2003) Targeted degradation of TOC1 by ZTL modulates circadian function in Arabidopsis thaliana. *Nature*, **426**, 567-570.
- Mathews S. & Sharrock R.A. (1997) Phytochrome gene diversity. *Plant, Cell & Environment*, **20**, 666-671.
- McClung C.R. & Davis S.J. (2010) Ambient Thermometers in Plants: From Physiological Outputs towards Mechanisms of Thermal Sensing. *Current Biology*, **20**, R1086-R1092.
- McWatters H.G., Bastow R.M., Hall A. & Millar A.J. (2000) The ELF3 zeitnehmer regulates light signalling to the circadian clock. *Nature*, **408**:716-720.
- McWatters H.G., Kolmos E., Hall A., Doyle M.R., Amasino R.M., Gyula P., Nagy F., Millar A.J. & Davis S.J. (2007) ELF4 Is Required for Oscillatory Properties of the Circadian Clock. *Plant Physiology*, **144**, 391-401.
- Michael T.P., Salomé P.A., Yu H.J., Spencer T.R., Sharp E.L., McPeck M.A., Alonso J.M., Ecker J.R. & McClung C.R. (2003) Enhanced Fitness Conferred by Naturally Occurring Variation in the Circadian Clock. *Science*, **302**, 1049-1053.
- Millar A., Straume M., Chory J., Chua N. & Kay S. (1995) The regulation of circadian period by phototransduction pathways in Arabidopsis. *Science*, **267**, 1163-1166.
- Millar A.J. & Kay S.A. (1996) Integration of circadian and phototransduction pathways in the network controlling CAB gene transcription in Arabidopsis. *Proceedings of the National Academy of Sciences*, **93**, 15491-15496.
- Millar A.J., McGrath R.B. & Chua N.H. (1994) PHYTOCHROME PHOTOTRANSDUCTION PATHWAYS. *Annual Review of Genetics*, **28**, 325-349.
- Mizoguchi T., Wright L., Fujiwara S., Cremer F., Lee K., Onouchi H., Mouradov A., Fowler S., Kamada H., Putterill J. & Coupland G. (2005) Distinct Roles of GIGANTEA in Promoting Flowering and Regulating Circadian Rhythms in Arabidopsis. *The Plant Cell*, **17**, 2255-2270.
- Mizuno T., Nomoto Y., Oka H., Kitayama M., Takeuchi A., Tsubouchi M. & Yamashino T. (2014) Ambient temperature signal feeds into the circadian clock transcriptional circuitry through the EC night-time repressor in Arabidopsis thaliana. *Plant and Cell Physiology*, **55**, 958-976.
- Mockler T., Michael T., Priest H., Shen R., Sullivan C., Givan S., McEntee C., Kay S. & Chory J. (2007) *The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis*. Paper presented at the Cold Spring Harbor Symposia on Quantitative Biology.
- Nakamichi N., Kita M., Ito S., Yamashino T. & Mizuno T. (2005) PSEUDO-RESPONSE REGULATORS, PRR9, PRR7 and PRR5, Together Play Essential Roles Close to the Circadian Clock of Arabidopsis thaliana. *Plant and Cell Physiology*, **46**, 686-698.
- Nakamichi N., Kusano M., Fukushima A., Kita M., Ito S., Yamashino T., Saito K., Sakakibara H. & Mizuno T. (2009) Transcript Profiling of an Arabidopsis PSEUDO RESPONSE REGULATOR Arrhythmic Triple Mutant Reveals a Role for the Circadian Clock in Cold Stress Response. *Plant and Cell Physiology*, **50**, 447-462.
- Nomoto Y., Kubozono S., Yamashino T., Nakamichi N. & Mizuno T. (2012) Circadian Clock- and PIF4-Controlled Plant Growth: A Coincidence Mechanism Directly Integrates a Hormone Signaling Network into the Photoperiodic Control of Plant Architectures in Arabidopsis thaliana. *Plant and Cell Physiology*, **53**, 1950-1964.

- 946 Nozue K., Covington M.F., Duek P.D., Lorrain S., Fankhauser C., Harmer S.L. & Maloof J.N. (2007)
 947 Rhythmic growth explained by coincidence between internal and external cues. *Nature*, **448**,
 948 358-361.
- 949 Nusinow D.A., Helfer A., Hamilton E.E., King J.J., Imaizumi T., Schultz T.F., Farre E.M. & Kay S.A.
 950 (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl
 951 growth. *Nature*, **475**, 398-402.
- 952 Onai K. & Ishiura M. (2005) PHYTOCLOCK 1 encoding a novel GARP protein essential for the
 953 Arabidopsis circadian clock. *Genes to Cells*, **10**, 963-972.
- 954 Oravecz A., Baumann A., Máté Z., Brzezinska A., Molinier J., Oakeley E.J., Ádám É., Schäfer E., Nagy F.
 955 & Ulm R. (2006) CONSTITUTIVELY PHOTOMORPHOGENIC1 Is Required for the UV-B
 956 Response in Arabidopsis. *The Plant Cell*, **18**, 1975-1990.
- 957 Ordoñez-Herrera N., Fackendahl P., Yu X., Schaefer S., Koncz C. & Hoecker U. (2015) A cop1 spa
 958 Mutant Deficient in COP1 and SPA Proteins Reveals Partial Co-Action of COP1 and SPA
 959 during Arabidopsis Post-Embryonic Development and Photomorphogenesis. *Molecular*
 960 *Plant*, **8**, 479-481.
- 961 Osterlund M.T., Ang L.H. & Deng X.W. (1999) The role of COP1 in repression of Arabidopsis
 962 photomorphogenic development. *Trends in Cell Biology*, **9**, 113-118.
- 963 Osterlund M.T., Hardtke C.S., Wei N. & Deng X.W. (2000) Targeted destabilization of HY5 during
 964 light-regulated development of Arabidopsis. *Nature*, **405**, 462-466.
- 965 Park D.H., Somers D.E., Kim Y.S., Choy Y.H., Lim H.K., Soh M.S., Kim H.J., Kay S.A. & Nam H.G. (1999)
 966 Control of Circadian Rhythms and Photoperiodic Flowering by the Arabidopsis GIGANTEA
 967 Gene. *Science*, **285**, 1579-1582.
- 968 Pokhilko A., Hodge S.K., Stratford K., Knox K., Edwards K.D., Thomson A.W., Mizuno T. & Millar A.J.
 969 (2010) Data assimilation constrains new connections and components in a complex,
 970 eukaryotic circadian clock model. *Molecular Systems Biology*, **6**.
- 971 Pyl E.T., Piques M., Ivakov A., Schulze W., Ishihara H., Stitt M. & Sulpice R. (2012) Metabolism and
 972 Growth in Arabidopsis Depend on the Daytime Temperature but Are Temperature-
 973 Compensated against Cool Nights. *The Plant Cell*, **24**, 2443-2469.
- 974 Qin M.M., Kuhn R., Moran S. & Quail P.H. (1997) Overexpressed phytochrome C has similar
 975 photosensory specificity to phytochrome B but a distinctive capacity to enhance primary leaf
 976 expansion. *Plant Journal*, **12**, 1163-1172.
- 977 Quail P., Boylan M., Parks B., Short T., Xu Y. & Wagner D. (1995) Phytochromes: photosensory
 978 perception and signal transduction. *Science*, **268**, 675-680.
- 979 Raschke A., Ibañez C., Ullrich K.K., Anwer M.U., Becker S., Glöckner A., Trenner J., Denk K., Saal B.,
 980 Sun X., Ni M., Davis S.J., Delker C. & Quint M. (2015) Natural variants of ELF3 affect
 981 thermomorphogenesis by transcriptionally modulating PIF4-dependent auxin response
 982 genes. *BMC Plant Biology*, **15**, 197.
- 983 Reed J.W., Nagpal P., Bastow R.M., Solomon K.S., Dowson-Day M.J., Elumalai R.P. & Millar A.J. (2000)
 984 Independent Action of ELF3 and phyB to Control Hypocotyl Elongation and Flowering Time.
 985 *Plant Physiology*, **122**, 1149-1160.
- 986 Rizzini L., Favory J.J., Cloix C., Faggionato D., O'Hara A., Kaiserli E., Baumeister R., Schäfer E., Nagy F.,
 987 Jenkins G.I. & Ulm R. (2011) Perception of UV-B by the Arabidopsis UVR8 Protein. *Science*,
 988 **332**, 103-106.
- 989 Roden L.C., Song H.-R., Jackson S., Morris K. & Carre I.A. (2002) Floral responses to photoperiod are
 990 correlated with the timing of rhythmic expression relative to dawn and dusk in Arabidopsis.
 991 *Proceedings of the National Academy of Sciences*, **99**, 13313-13318.
- 992 Rudiger W., Thummler F., Cmiel E. & Schneider S. (1983) Chromophore Structure of the
 993 Physiologically Active Form (Pfr) of Phytochrome. *Proceedings of the National Academy of*
 994 *Sciences of the United States of America-Biological Sciences*, **80**, 6244-6248.
- 995 Sakamoto K. & Nagatani A. (1996) Nuclear localization activity of phytochrome B. *The Plant Journal*,
 996 **10**, 859-868.

Formatted: Font: Calibri, 11 pt, Not Bold,
Font color: Auto

Formatted: Font: Calibri, 11 pt, Font color:
Auto

Formatted: Font: Calibri, 11 pt, Not Bold,
Font color: Auto

Formatted: Font: Calibri, 11 pt, Not Bold,
Font color: Auto

- 997 Salazar J.D., Saithong T., Brown P.E., Foreman J., Locke J.C.W., Halliday K.J., Carré I.A., Rand D.A. &
998 Millar A.J. (2009) Prediction of Photoperiodic Regulators from Quantitative Gene Circuit
999 Models. *Cell*, **139**, 1170-1179.
- 1000 Salomé P.A., Michael T.P., Kearns E.V., Fett-Neto A.G., Sharrock R.A. & McClung C.R. (2002) The out
1001 of phase 1 Mutant Defines a Role for PHYB in Circadian Phase Control in Arabidopsis. *Plant*
1002 *Physiology*, **129**, 1674-1685.
- 1003 Salter M.G., Franklin K.A. & Whitelam G.C. (2003) Gating of the rapid shade-avoidance response by
1004 the circadian clock in plants. *Nature*, **426**, 680-683.
- 1005 Sanchez A., Shin J. & Davis S.J. (2011) Abiotic stress and the plant circadian clock. *Plant Signaling &*
1006 *Behavior*, **6**, 223-231.
- 1007 Sanchez-Villarreal A., Shin J., Bujdosó N., Obata T., Neumann U., Du S., Ding Z., Davis A.M., Shindo T.,
1008 Schmelzer E., Sulpice R., Nunes-Nesi A., Stitt M., Fernie A.R. & Davis S.J. (2013) TIME FOR
1009 COFFEE is an essential component in the maintenance of metabolic homeostasis in
1010 Arabidopsis thaliana. *The Plant Journal*, **76**, 188-200.
- 1011 Schaffer R., Ramsay N., Samach A., Corden S., Putterill J., Carré I.A. & Coupland G. (1998) The late
1012 elongated hypocotyl Mutation of Arabidopsis Disrupts Circadian Rhythms and the
1013 Photoperiodic Control of Flowering. *Cell*, **93**, 1219-1229.
- 1014 Sellaro R., Pacín M. & Casal J.J. (2012) Diurnal Dependence of Growth Responses to Shade in
1015 Arabidopsis: Role of Hormone, Clock, and Light Signaling. *Molecular Plant*, **5**, 619-628.
- 1016 Seo H.S., Yang J.Y., Ishikawa M., Bolle C., Ballesteros M.L. & Chua N.H. (2003) LAF1 ubiquitination by
1017 COP1 controls photomorphogenesis and is stimulated by SPA1. *Nature*, **423**, 995-999.
- 1018 Sharrock R.A. & Quail P.H. (1989) Novel phytochrome sequences in Arabidopsis thaliana: structure,
1019 evolution, and differential expression of a plant regulatory photoreceptor family. *Genes &*
1020 *Development*, **3**, 1745-1757.
- 1021 Shearman L.P., Sriram S., Weaver D.R., Maywood E.S., Chaves I., Zheng B., Kume K., Lee C.C., van der
1022 G.T.J., Horst, Hastings M.H. & Reppert S.M. (2000) Interacting Molecular Loops in the
1023 Mammalian Circadian Clock. *Science*, **288**, 1013-1019.
- 1024 Sheerin D.J., Menon C., zur Oven-Krockhaus S., Enderle B., Zhu L., Johnen P., Schleifenbaum F.,
1025 Stierhof Y.D., Huq E. & Hiltbrunner A. (2015) Light-Activated Phytochrome A and B Interact
1026 with Members of the SPA Family to Promote Photomorphogenesis in Arabidopsis by
1027 Reorganizing the COP1/SPA Complex. *The Plant Cell*, **27**, 189-201.
- 1028 Shin J., Anwer M.U. & Davis S.J. (2013) Phytochrome-Interacting Factors (PIFs) as Bridges between
1029 Environmental Signals and the Circadian Clock: Diurnal Regulation of Growth and
1030 Development. *Molecular Plant*, **6**, 592-595.
- 1031 Shin J., Heidrich K., Sanchez-Villarreal A., Parker J.E. & Davis S.J. (2012) TIME FOR COFFEE Represses
1032 Accumulation of the MYC2 Transcription Factor to Provide Time-of-Day Regulation of
1033 Jasmonate Signaling in Arabidopsis. *Plant Cell*, **24**, 2470-2482.
- 1034 Shin J., Sánchez-Villarreal A., Davis A.M., Du S., Berendzen K.W., Koncz C., Ding Z., Li C. & Davis S.J.
1035 (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-
1036 dependent manner. *Plant, Cell & Environment*, **40**, 997-1008.
- 1037 Smith H. (1995) Physiological and Ecological Function within the Phytochrome Family. *Annual Review*
1038 *of Plant Physiology and Plant Molecular Biology*, **46**, 289-315.
- 1039 Somers D.E., Devlin P.F. & Kay S.A. (1998a) Phytochromes and Cryptochromes in the Entrainment of
1040 the Arabidopsis Circadian Clock. *Science*, **282**, 1488-1490.
- 1041 Somers D.E., Webb A.A., Pearson M. & Kay S.A. (1998b) The short-period mutant, *toc1-1*, alters
1042 circadian clock regulation of multiple outputs throughout development in Arabidopsis
1043 thaliana. *Development*, **125**, 485-494.
- 1044 Song Y.H., Smith R.W., To B.J., Millar A.J. & Imaizumi T. (2012) FKF1 Conveys Timing Information for
1045 CONSTANS Stabilization in Photoperiodic Flowering. *Science*, **336**, 1045-1049.

- 1046 Soy J., Leivar P., González-Schain N., Sentandreu M., Prat S., Quail P.H. & Monte E. (2012)
 1047 Phytochrome-imposed oscillations in PIF3 protein abundance regulate hypocotyl growth
 1048 under diurnal light/dark conditions in Arabidopsis. *The Plant Journal*, **71**, 390-401.
- 1049 [Soy J., Leivar P., González-Schain N., Martin G., Diaz C., Sentandreu M., Al-Sady B., Quail P. H.,
 1050 Monte E. \(2016\) Molecular convergence of clock and photosensory pathways through PIF3–
 1051 TOC1 interaction and co-occupancy of target promoters. *Proceedings of the National
 1052 Academy of Sciences of the United States of America*, **113** 4870-4875.](#)
- 1053 Stracke R., Favory J.J., Gruber H., Bartelniewoehner L., Bartels S., Binkert M., Funk M., Weisshaar B.
 1054 & Ulm R. (2010) The Arabidopsis bZIP transcription factor HY5 regulates expression of the
 1055 PFG1/MYB12 gene in response to light and ultraviolet-B radiation. *Plant, Cell & Environment*,
 1056 **33**, 88-103.
- 1057 Strayer C., Oyama T., Schultz T.F., Raman R., Somers D.E., Más P., Panda S., Kreps J.A. & Kay S.A.
 1058 (2000) Cloning of the Arabidopsis Clock Gene TOC1, an Autoregulatory Response Regulator
 1059 Homolog. *Science*, **289**, 768-771.
- 1060 Takase M., Mizoguchi T., Kozuka T. & Tsukaya H. (2013) The unique function of the Arabidopsis
 1061 circadian clock gene PRR5 in the regulation of shade avoidance response. *Plant Signaling &
 1062 Behavior*, **8**, e23534.
- 1063 Thines B. & Harmon F.G. (2010) Ambient temperature response establishes ELF3 as a required
 1064 component of the core Arabidopsis circadian clock. *Proceedings of the National Academy of
 1065 Sciences of the United States of America*, **107**, 3257-3262.
- 1066 Toth R., Kevei E., Hall A., Millar A.J., Nagy F. & Kozma-Bognár L. (2001) Circadian clock-regulated
 1067 expression of phytochrome and cryptochrome genes in Arabidopsis. *Plant Physiology*, **127**,
 1068 1607-1616.
- 1069 Tóth R., Kevei É., Hall A., Millar A.J., Nagy F. & Kozma-Bognár L. (2001) Circadian Clock-Regulated
 1070 Expression of Phytochrome and Cryptochrome Genes in Arabidopsis. *Plant Physiology*, **127**,
 1071 1607-1616.
- 1072 Troein C., Locke J.C.W., Turner M.S. & Millar A.J. (2009) Weather and Seasons Together Demand
 1073 Complex Biological Clocks. *Current Biology*, **19**, 1961-1964.
- 1074 Undurraga S.F., Press M.O., Legendre M., Bujdosó N., Bale J., Wang H., Davis S.J., Verstrepen K.J. &
 1075 Queitsch C. (2012) Background-dependent effects of polyglutamine variation in the
 1076 Arabidopsis thaliana gene ELF3. *Proceedings of the National Academy of Sciences of the
 1077 United States of America*, **109**, 19363-19367.
- 1078 Walters R.G., Rogers J.J.M., Shephard F. & Horton P. (1999) Acclimation of Arabidopsis thaliana to
 1079 the light environment: the role of photoreceptors. *Planta*, **209**, 517-527.
- 1080 Wang H., Ma L.G., Li J.M., Zhao H.Y. & Deng X.W. (2001) Direct interaction of Arabidopsis
 1081 Cryptochromes with COP1 in Light Control Development. *Science*, **294**, 154-158.
- 1082 Wang W., Barnaby J.Y., Tada Y., Li H., Tor M., Caldelari D., Lee D.U., Fu X.D. & Dong X. (2011) Timing
 1083 of plant immune responses by a central circadian regulator. *Nature*, **470**, 110-114.
- 1084 Wang Z.Y. & Tobin E.M. (1998) Constitutive Expression of the CIRCADIAN CLOCK ASSOCIATED 1
 1085 (CCA1) Gene Disrupts Circadian Rhythms and Suppresses Its Own Expression. *Cell*, **93**, 1207-
 1086 1217.
- 1087 Wenden B., Kozma-Bognár L., Edwards K.D., Hall A.J., Locke J.C. & Millar A.J. (2011) Light inputs
 1088 shape the Arabidopsis circadian system. *Plant J*, **66**, 480-491.
- 1089 Weston E., Thorogood K., Vinti G. & López-Juez E. (2000) Light quantity controls leaf-cell and
 1090 chloroplast development in Arabidopsis thaliana wild type and blue-light-perception
 1091 mutants. *Planta*, **211**, 807-815.
- 1092 Yang H.Q., Wu Y.J., Tang R.H., Liu D., Liu Y. & Cashmore A.R. (2000) The C Termini of Arabidopsis
 1093 Cryptochromes Mediate a Constitutive Light Response. *Cell*, **103**, 815-827.
- 1094 Yang J., Lin R., Sullivan J., Hoecker U., Liu B., Xu L., Deng X.W. & Wang H. (2005) Light Regulates
 1095 COP1-Mediated Degradation of HFR1, a Transcription Factor Essential for Light Signaling in
 1096 Arabidopsis. *The Plant Cell*, **17**, 804-821.

Formatted: English (U.S.), Do not check spelling or grammar

Formatted: Font: Calibri, 11 pt, Not Bold, Font color: Text 1

Formatted: Font: Calibri, 11 pt, Font color: Text 1

Formatted: Font: Calibri, 11 pt, Not Bold, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Calibri, 11 pt, Not Bold, Font color: Text 1

1097 Yanovsky M.J., Mazzella M.A. & Casal J.J. (2000) A quadruple photoreceptor mutant still keeps track
1098 of time. *Current Biology*, **10**, 1013-1015.

1099 Yi C. & Deng X.W. (2005) COP1 – from plant photomorphogenesis to mammalian tumorigenesis.
1100 *Trends in Cell Biology*, **15**, 618-625.


1101 Yu J.W., Rubio V., Lee N.Y., Bai S., Lee S.Y., Kim S.S., Liu L., Zhang Y., Irigoyen M.L., Sullivan J.A., Zhang
1102 Y., Lee I., Xie Q., Paek N.C. & Deng X.W. (2008) COP1 and ELF3 Control Circadian Function
1103 and Photoperiodic Flowering by Regulating GI Stability. *Molecular Cell*, **32**, 617-630.

1104 Zagotta M.T., Hicks K.A., Jacobs C.I., Young J.C., Hangarter R.P. & MeeksWagner D.R. (1996) The
1105 Arabidopsis ELF3 gene regulates vegetative photomorphogenesis and the photoperiodic
1106 induction of flowering. *Plant Journal*, **10**, 691-702.

1107 Zeilinger M.N., Farré E.M., Taylor S.R., Kay S.A. & Doyle F.J. (2006) A novel computational model of
1108 the circadian clock in Arabidopsis that incorporates PRR7 and PRR9. *Molecular Systems
1109 Biology*, **2**.

1110 Zhu D., Maier A., Lee J.H., Laubinger S., Saijo Y., Wang H., Qu L.J., Hoecker U. & Deng X.W. (2008)
1111 Biochemical Characterization of Arabidopsis Complexes Containing CONSTITUTIVELY
1112 PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA Proteins in Light Control of Plant
1113 Development. *The Plant Cell*, **20**, 2307-2323.

1114 Zuo Z., Liu H., Liu B., Liu X. & Lin C. (2011) Blue Light-Dependent Interaction of CRY2 with SPA1
1115 Regulates COP1 activity and Floral Initiation in Arabidopsis. *Current biology : CB*, **21**, 841-847.



ELF3 localisation: a master integrator of light-input to the circadian clock

361x361mm (72 x 72 DPI)

